



CME POST-TEST

All post-tests must be completed and submitted online.

EXPIRATION DATE: NOVEMBER 2023

Earn Category I CME Credit by reading both CME articles in this issue, reviewing the post-test, then taking the online test at <http://cme.aapa.org>. Successful completion is defined as a cumulative score of at least 70% correct. This material has been reviewed and is approved for 1 hour of clinical Category I (Preapproved) CME credit by the AAPA.

PANCREATOGENIC DIABETES

1. **Type 3c diabetes is also known as**
 - a. type 1 diabetes
 - b. type 2 diabetes
 - c. pancreatogenic diabetes
 - d. insulin-dependent diabetes
2. **Chronic pancreatitis accounts for what percentage of cases of type 3c diabetes?**
 - a. 61%
 - b. 67%
 - c. 70%
 - d. 79%
3. **Pancreatic exocrine insufficiency is near-universal in patients with**
 - a. liver cirrhosis
 - b. morbid obesity
 - c. adrenal adenoma
 - d. cystic fibrosis
4. **According to the ADA, in addition to standard criteria for diagnosing diabetes, diagnosis of type 3c diabetes requires**
 - a. presence of pancreatic insufficiency, evidence of pathologic pancreatic imaging, and absence of type 1 diabetes-associated autoimmune markers
 - b. elevated liver enzymes, evidence of pathologic pancreatic imaging, and absence of type 1 diabetes-associated autoimmune markers
 - c. presence of pancreatic insufficiency, a history of chronic pancreatitis, and absence of type 1 diabetes-associated autoimmune markers
 - d. presence of pancreatic insufficiency, evidence of pathologic pancreatic imaging, and a history of ketoacidosis
5. **The most often recommended therapy for patients with type 3c diabetes is**
 - a. glipizide
 - b. rosiglitazone
 - c. insulin
 - d. semaglutide

BLUNT CHEST WALL TRAUMA

6. **Injuries associated with blunt chest wall trauma have a mortality of**
 - a. 2% to 5%
 - b. 5% to 10%
 - c. 10% to 20%
 - d. 20% to 30%
7. **Rib fractures account for what percent of blunt traumatic injuries in adults?**
 - a. 25%
 - b. 35%
 - c. 45%
 - d. 55%
8. **Which imaging study is the standard of care for evaluation in hemodynamically stable trauma patients with chest wall injury?**
 - a. MRI
 - b. CT imaging
 - c. chest radiograph
 - d. ultrasound
9. **The Western Trauma Association recommends ICU admission for**
 - a. all patients with more than two rib fractures
 - b. patients over age 65 years with more than three significant rib fractures
 - c. patients over age 70 years with more than two significant rib fractures
 - d. patients over age 65 years with more than two significant rib fractures
10. **In which patients should surgical stabilization be considered?**
 - a. all patients with a flail chest or with three or more severely displaced ribs
 - b. all patients with a flail chest and with five or more severely displaced ribs
 - c. only patients over age 65 years with a flail chest or with three or more severely displaced ribs
 - d. all patients with traumatic tension pneumothorax and three or more severely displaced ribs