



# Practical Strategies for Engaging Individuals With Obesity in Primary Care

Scott I. Kahan, MD, MPH

## Abstract

Although widely recognized as a chronic disease that requires long-term, structured, and multidisciplinary management, obesity remains largely underdiagnosed and undertreated. The prevalence of obesity continues to increase dramatically, with the highest rates seen in the United States. Despite the availability of several clinical practice guidelines, published studies suggest that health care professionals (HCPs) infrequently and inconsistently follow guideline recommendations. Barriers to HCP participation in obesity management are likely to inhibit obesity counseling in primary care. Improving HCP obesity-related practices and counseling is important. This article discusses current practices, barriers to effective obesity management, and recommendations to improve HCP obesity management and counseling, based on findings from a PubMed search and clinical experience. The aim of the article is to share best-practice strategies for engaging patients.

© 2018 Mayo Foundation for Medical Education and Research. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>) ■ *Mayo Clin Proc.* 2018;93(3):351-359

The past half-century has seen a staggering increase in global obesity rates, with 266 million men and 375 million women now exceeding the weight threshold for obesity (defined by a body mass index [BMI; calculated as weight in kilograms divided by height in meters squared] of  $\geq 30$  kg/m<sup>2</sup>).<sup>1</sup> The prevalence of obesity (BMI, 30-40 kg/m<sup>2</sup>) and severe obesity (BMI,  $\geq 40$  kg/m<sup>2</sup>) is approximately 37.7% and 7.7%, respectively,<sup>2</sup> representing an approximate increase of 10% in the prevalence of obesity in the United States since 2005–2006 and an increase in excess of 600% in the prevalence of severe obesity since the mid-1980s.<sup>2,3</sup> In addition to being a public health epidemic affecting large populations, obesity is a complex and multifaceted clinical condition. Several global organizations and regulatory bodies explicitly recognize obesity as a chronic disease, including the World Health Organization, the US Food and Drug Administration, the US National Institutes of Health, and, most recently, the American Medical Association.<sup>4-8</sup> Obesity is associated with almost 200 metabolic, mechanical, and mental comorbid conditions,<sup>9</sup> the prevalence of which increase as BMI rises.<sup>10</sup> People with obesity have an increased risk of development of cardiovascular disease and type 2 diabetes<sup>11</sup> and a reduced

health-related quality of life.<sup>12</sup> The financial costs of obesity are substantial, both to affected individuals and society at large.<sup>13</sup>

As with other chronic conditions, successful management is not necessarily defined by achieving a “cure.” Long-term weight loss is generally accepted as maintaining weight loss of 5% to 10% for 1 or more years and is associated with improvements in major risk factors.<sup>14-21</sup> Conventional wisdom and older research suggest that long-term weight loss success is elusive<sup>22,23</sup>; however, recent studies have shown impressive outcomes for behavioral weight loss interventions, particularly when provided by trained interventionists. The Look AHEAD study evaluated weight loss achieved with behavioral weight loss counseling in 5145 people with overweight or obesity and type 2 diabetes.<sup>24</sup> Patients receiving behavioral counseling lost more than 8% of body weight during the first year and, at year 8, 50.0% of intervention patients maintained a 5% or more loss, 26.9% maintained a greater than 10% loss, and 11.0% maintained a more than 15% loss of initial body weight. Systematic reviews of behavioral weight loss counseling conducted by the US Preventive Services Task Force (USPSTF) and the National Heart, Lung, and Blood Institute (NHLBI) Obesity Expert Panel found similarly

From Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, and George Washington University School of Medicine and Health Sciences, Washington, DC.

## ARTICLE HIGHLIGHTS

- Evidence indicates that health care professional (HCP)—patient engagement on the topic of obesity is often suboptimal.
- Lack of HCP—patient engagement can adversely affect the diagnosis and management of obesity.
- Barriers perceived by HCPs and patients influence their level of engagement.
- Several strategies may help to address the barriers and enable effective HCP—patient engagement.
- The health benefits that patients derive from being engaged with their HCP are clearly evident.

impressive outcomes, which has led to policy changes encouraging behavioral counseling in primary care.<sup>20,25</sup> Primary care physicians (PCPs) are in the unique position of engaging people with obesity and ensuring that they receive appropriate counseling. Nonetheless, whereas most people with obesity have tried—often repeatedly—to lose weight,<sup>26</sup> few receive formal guidance and counseling from their health care professional (HCP).<sup>27</sup> Based on relevant articles from a PubMed search performed in June 2017 and my own clinical experience, this article discusses current obesity-related practices of HCPs in the United States and the availability of specialist training for US HCPs and provides practical recommendations for improving how HCPs engage with patients to support long-term weight management.

## CURRENT GUIDELINES AND PRACTICE

**The Importance of Screening and Diagnosis**

Screening and diagnosis are fundamental steps in addressing medical conditions. In a study of 7790 National Health and Nutrition Examination Survey (NHANES) participants, those whose weight problem was diagnosed and who were advised of their weight status by their HCP had 6.1-fold (BMI, >25 kg/m<sup>2</sup>) or 8.3-fold (BMI, >30 kg/m<sup>2</sup>) increased odds of correctly perceiving their weight status.<sup>28</sup> Moreover, participants had 2.5-fold (BMI, >25 kg/m<sup>2</sup>) or 2.2-fold (BMI, >30 kg/m<sup>2</sup>) increased odds of attempting to lose weight. Among 9827 patients in a large primary care

database, formal diagnosis of obesity was associated with 2.4-fold increased odds of having an obesity management plan in place.<sup>29</sup> An NHANES analysis of 5054 participants revealed 1.9-fold (BMI, >25 kg/m<sup>2</sup>) and 1.8-fold (BMI, >30 kg/m<sup>2</sup>) increased odds of losing weight (defined as reported >5% weight loss) if their HCP had diagnosed and communicated that they were overweight.<sup>30</sup>

**Current Guidelines for Screening and Diagnosis**

The USPSTF recommends that clinicians screen all adults for obesity and offer referral for intensive multicomponent behavioral interventions to patients with a BMI of 30 kg/m<sup>2</sup> or higher.<sup>31</sup> The NHLBI obesity guidelines, disseminated by the American Heart Association/American College of Cardiology/The Obesity Society, recommend that weight should be measured and BMI calculated at least every year in all people who need to lose weight.<sup>32</sup> Body mass index screening and follow-up is a clinical quality measure now included in the Physician Quality Reporting System<sup>33</sup> and carried out by the Centers for Medicare and Medicaid Services.<sup>34</sup> Moreover, negative payment adjustments are levied to HCPs and group practices who do not satisfactorily report data on quality measures for Medicare Part B Physician Fee Schedule-covered professional services.<sup>34</sup>

**Current Practice for Screening and Diagnosis**

Despite several guidelines, practice recommendations, and incentives, these measures are not consistently followed in primary care settings.<sup>35</sup> According to retrospective analyses of patient records, up to 90% of people with obesity (BMI, ≥30 to ≤35 kg/m<sup>2</sup>) have not received a formal diagnosis from their physician.<sup>36</sup> Although rates of diagnosis increase in patients in higher weight classes, they remain suboptimal across all weights.<sup>35</sup>

**The Importance of Obesity Counseling**

Counseling delivered by HCPs substantially affects weight loss behaviors. A cross-sectional analysis of 810 British adults with a BMI of less than 25 kg/m<sup>2</sup> revealed that an HCP recommendation to lose weight was associated with 3.7-fold increased odds of wanting to weigh less and 3.5-fold increased odds of attempting to

lose weight.<sup>37</sup> A 2013 meta-analysis of 32 published studies on the impact of HCP weight loss counseling on actual changes in patient behavior found that patients are nearly 4 times more likely to attempt weight loss when their HCP provides weight loss advice.<sup>38</sup>

### Current Guidelines for Obesity Counseling

All relevant guidelines, including the USPSTF and the NHLBI-led American Heart Association/American College of Cardiology/The Obesity Society recommendations, highlight intensive behavioral counseling for patients with a BMI of 30 kg/m<sup>2</sup> or higher.<sup>2,20,32</sup> Counseling should be intensive initially (usually defined by weekly or biweekly interactions) followed by monthly interactions (after 6 months), should be multicomponent, and should involve guidance for long-term maintenance. Beginning in 2011, Medicare began coverage for intensive obesity counseling in primary care, consisting of weekly obesity counseling visits for 1 month, followed by biweekly visits for months 2 through 6, followed by monthly visits thereafter.<sup>39</sup>

### Current Practice for Obesity Counseling

Despite these guidelines and coverage, only a minority of HCPs engage in weight loss counseling with their patients. In 2008, a survey of more than 1200 PCPs sampled from the American Medical Association Masterfile found that less than 50% of PCPs reported always providing advice on diet, exercise, or weight control.<sup>40</sup> A cross-sectional study of 21,220 US adult outpatient visits with 954 PCPs between 2007 and 2008 (taken from the National Ambulatory Medical Care Survey [NAMCS] data from 32,500 adult primary care visits to PCPs) revealed that 58% of PCPs performed no weight counseling during clinic appointments.<sup>41</sup> Only 85 physicians in the NAMCS survey (8.9%) provided the majority of the weight counseling, suggesting that the vast majority of PCPs perform little or no weight counseling.<sup>41</sup> The NAMCS survey data for 2012 show that just 4.9% of adult office visits in the previous year were for obesity, despite obesity prevalence approaching 40% in the US.<sup>42</sup>

Some studies suggest that the rates of weight loss counseling in primary care may actually be declining. Among 10,113 primary

care visits in 2003–2004, fewer than 5.8% included weight loss counseling, which represented a 30% decline from 1995 through 1996.<sup>43</sup> Data from the NAMCS survey revealed that weight loss counseling declined from 7.8% of visits in 1995–1996 to 6.2% of visits in 2007–2008, despite the increasing prevalence of obesity. A study that evaluated secular trends in the diagnosis and treatment of obesity among an NHANES sample of 31,039 nonpregnant adults found no change in the odds of receiving a diagnosis of obesity from a physician or physician-directed weight loss from 1994 through 2008, despite a large increase (from 56.1% to 69.1%) in the prevalence of overweight in US adults during that period.<sup>44</sup> Furthermore, several studies report minimal knowledge of, and adherence to, published guidelines for obesity counseling and treatment among PCPs.<sup>45,46</sup>

In terms of the patient perspective on current practice for obesity counseling, a 2014 study of 1002 male firefighters reported that 69% of participants overall and 48% of participants with obesity reported never having received weight loss advice from their HCPs.<sup>47</sup> Although the likelihood of receiving weight loss advice increased with age and severity of obesity, data from the same study suggest that just 68.1% of firefighters with class II/III obesity received weight loss advice from their HCP.<sup>47</sup> Among 2725 US adults with a mean BMI of 37.1 kg/m<sup>2</sup>, almost one-quarter responded that they had never discussed their weight with their PCP, and nearly two-thirds reported that they wished their physician would discuss weight or offer weight management counseling to them.<sup>48</sup> Furthermore, this problem is not limited to the US health care system. For example, the aforementioned 2013 cross-sectional analysis of 810 British adults with a BMI higher than 25 kg/m<sup>2</sup> reported that just 17% and 42% of people with overweight or obesity, respectively, recalled ever receiving advice to lose weight from their HCP.<sup>37</sup>

### Perceived Barriers

Health care professionals perceive several barriers that prevent them from fully aligning their clinical practice with current guideline recommendations. Lack of training in obesity is a key barrier. Just 25% of medical schools

**TABLE. Considerations for the Clinic Environment**

Furniture	<ul style="list-style-type: none"> <li>● Wide-base, higher-weight-capacity chairs (preferably armless) in both the waiting and patient areas</li> <li>● Provide specialized bariatric chairs, where possible</li> <li>● Pedestal toilets, rather than wall-mounted</li> <li>● Wheelchair-accessible bathrooms</li> </ul>
Equipment	<ul style="list-style-type: none"> <li>● Large-sized or thigh-sized blood pressure cuffs</li> <li>● High-capacity weighing scales (ideally &gt;500 lbs [to convert to kg, multiply by 0.45])</li> <li>● Extra-large gowns</li> </ul>
Staff	<ul style="list-style-type: none"> <li>● Educated about obesity and weight bias</li> </ul>

Modified from the Strategies to Overcome and Prevent (STOP) Obesity Alliance website.<sup>41</sup>

offer a dedicated nutrition course—a decline since the status of nutrition education in US medical schools was first assessed in 1985—and few medical schools achieve the 25 hours of nutrition education recommended by the National Academy of Sciences.<sup>49</sup> A review of obesity-related content on the US Medical Licensing Examinations found that obesity-specific content was rarely included in examinations and most relevant topics were entirely absent.<sup>50</sup> Additional barriers often cited by HCPs as a reason for not following counseling guideline recommendations include insufficient training and counseling skills,<sup>51-53</sup> limited time,<sup>53-56</sup> lack of confidence that obesity counseling is worthwhile,<sup>52</sup> shortage of appropriate clinical space and resources,<sup>52,56-59</sup> and the challenges of reimbursement.<sup>54,58</sup> These barriers likely result in missed opportunities to engage patients in discussions about weight, to screen for obesity, and to engage in weight loss counseling.

From a patient perspective, barriers to effective screening and diagnosis may occur even before they have visited their HCP. Many people with overweight or obesity feel stigmatized and unwelcome in the clinic (eg, waiting room furniture, weighing scales, or examination tables are of insufficient capacity for heavier patients), have had prior negative health care experiences (such as being shamed by physicians to lose weight), or believe that seeking help signifies moral failure.<sup>51,57</sup> The stigma of having a formal diagnosis of obesity may also prevent individuals from consulting a physician, and evidence suggests that patients with higher BMIs experience

less respect from physicians.<sup>60,61</sup> Additionally, patients often expect HCPs to proactively broach the topic of excess weight; if weight is not discussed by HCPs, patients may assume it is not a concern.<sup>62</sup> Even when obesity is discussed, in many cases no follow-up appointment for ongoing care is scheduled.<sup>55</sup> These barriers may result in patients delaying preventive screening, which results in increased health risks associated with obesity because diagnosis is delayed.<sup>57</sup>

## PRACTICAL GUIDANCE FOR CLINICAL PRACTICE

Several relatively small considerations can improve HCP–patient interaction regarding obesity and improve obesity-related care and outcomes.

### Using People-First Language and Motivating Terminology

The words used to describe excess weight can be polarizing and may be demotivating. For example, the term *obese* is consistently rated as least desirable and off-putting by patients.<sup>63,64</sup> In contrast, neutral terms, such as *unhealthy weight*, are generally preferred and may be perceived as motivating, thereby helping patients feel supported and empowered in their weight management efforts.<sup>63,64</sup> As with other medical conditions, the use of people-first language (“patient with obesity” or “having obesity”), as opposed to condition-first language (“obese patient”), is recommended terminology for both academic and clinical settings.<sup>65,66</sup>

### Ensuring an Appropriate Clinic Environment

Paying attention to the physical requirements of the clinic space and procedures can greatly improve the health care experience for patients (Table) and is more likely to encourage them to seek help and to be receptive to counseling. Even simple steps can be beneficial, such as ensuring that scales are in private areas and offering medical equipment that is suitable for larger individuals.<sup>67</sup>

### Integrating Digital Technology

Technological aids—such as electronic reminders to prompt BMI measurement, which are often standard in electronic health record (EHR) programs—can considerably improve obesity diagnosis and documentation. In a recent

study in which physicians were randomized to receive a BMI reminder prompt as part of patients' EHR, they were 3 times more likely to document a diagnosis of overweight during a patient consultation than physicians who did not receive the reminder (22% vs 7%;  $P=.02$ ).<sup>68</sup> Notably, 91% of physicians who recorded BMI using the EHR admitted the tool alerted them to patients they did not realize were overweight.


### Use a Structured Framework for Counseling

Numerous frameworks are available to help HCPs structure obesity counseling. A useful framework for counseling is the "5As" (Ask, Assess, Advise, Agree, and Assist) intervention strategy (Figure), which was initially developed to support smoking cessation and has been adapted for obesity management.<sup>70</sup>

A 2014 study found that implementing the 5As framework in primary care clinics doubled the initiation of obesity management compared with clinics that did not use the intervention, was easy to implement, and resulted in positive health-related behavioral changes by patients.<sup>71</sup> A 2013 study that randomized internal medicine residents to receive structured training in the 5As via an obesity counseling curriculum vs receiving standard residency training reported that patients attending appointments with the 5As-trained residents lost significantly more weight over the course of 12 months compared with patients of the residents in the control group, who gained weight.<sup>72</sup> A study conducted in 2010 found that patients who received counseling using parts of the 5As framework had more than 30% greater odds of being motivated to lose weight.<sup>73</sup>

### Begin With Small Steps

Analogous to the advice that patients receive regarding the value of small steps for behavioral change, HCPs should follow the same advice. Simply starting a dialogue about obesity by asking the patient's permission to discuss weight, as described in the 5As, may help to broach an often sensitive topic and is less likely to be interpreted as judgmental.<sup>70,71</sup> Asking permission conveys empathy and respect for the patient, which has been shown to increase motivation for behavioral changes in nutrition and physical activity, whereas



**FIGURE.** The 5As framework for obesity counseling. BMI = body mass index; SMART = smart, measurable, achievable, relevant, time-based; WC = waist circumference; WL = weight loss. Modified from the Canadian Obesity Network website.<sup>69</sup>

perceived judgement about weight negatively influences weight loss.<sup>74</sup>

A recent randomized controlled trial in which physicians were trained to engage in a 30-second obesity intervention found that even exceedingly small steps can lead to positive outcomes.<sup>75</sup> Physicians were randomized to an intervention group in which they initiated a discussion regarding weight, offered patients referral to a weight management program, and offered a follow-up appointment compared with a control group that only advised patients that weight loss would improve their health. Both physicians and patients found the intervention acceptable and helpful. Patients counseled by physicians in the intervention group lost more than twice as much weight over 1 year and had more than twice the odds of losing at least 5% and at least 10% of their starting body weight.

### Use Available Educational Resources

For practicing HCPs who do not have the benefit of ongoing improvements in obesity

training in medical schools, it is now becoming easier for them to improve their knowledge about obesity via self-study or in-person or online continuing medical education. Many professional societies also offer tools, guidance, and continuing education on obesity in order to improve HCP knowledge of and confidence in screening for and diagnosing obesity.<sup>58,76,77</sup> In 2012, the American Board of Obesity Medicine began certifying physicians who specialize in obesity to become “diplomates.”<sup>78</sup> Because of the lack of obesity medicine fellowship options, American Board of Obesity Medicine certification requires extensive continuing medical education credits, which has led to growing options for this type of HCP education.<sup>78</sup> A recent systematic review found that training physicians to improve communication and counseling regarding obesity leads to improved patient outcomes, including lowered BMI.<sup>79</sup>

### Do Not Go It Alone

Training in counseling techniques, such as motivational interviewing, is challenging for many HCPs.<sup>62</sup> However, a multidisciplinary team approach can minimize the burden of obesity management by sharing the patient’s care across HCPs, both within and outside the clinic.<sup>58</sup> Multidisciplinary team counseling approaches also help people maintain long-term weight loss. Employment of physician extenders, such as a dietitian, health coach, behavioral counselor, or other HCP who is trained in obesity management, should be considered.<sup>80,81</sup> In one study that evaluated the effectiveness of on-site dietitians to provide obesity and lipid counseling at an outpatient physician office, 80 patients seen by a dietitian at 2 visits achieved an average maximum weight loss of 5.6% at a mean follow-up of 1.75 years. It was anticipated that earlier and more frequent follow-up could result in greater and quicker weight loss.<sup>82</sup> Further, integrating community resources for obesity management with clinical care can significantly extend the reach and impact of HCPs with respect to obesity counseling.<sup>83</sup>

### Use Available Reimbursement Options

Although insufficient reimbursement is cited frequently as a key barrier to effective obesity counseling, access and coverage of counseling

is steadily improving. For example, in 2011, Medicare began reimbursing for intensive obesity counseling in primary care.<sup>84</sup> However, existing reimbursement options are rarely used by HCPs. A recent study evaluating the use of the Medicare reimbursement for obesity counseling reported that fewer than 1% of eligible patients received this counseling.<sup>85</sup> Both Medicaid and private health insurers are increasing their coverage for obesity counseling and related services, and this trend is likely to continue in the future. Health care professionals should attempt to maximize their use of reimbursed services.

### Long-term Support Is Required

Many patients do not adhere to sufficient long-term weight management support from HCPs.<sup>80,86</sup> Extended-duration weight loss counseling significantly improves long-term weight loss and maintenance of weight loss.<sup>87</sup> More intensive counseling over the long term (up to 5 face-to-face counseling sessions with a physical activity specialist and a registered dietitian over a 6-month period) has been reported to result in greater patient motivation and a greater reduction in weight, blood pressure, and cholesterol level compared with patients who only received standard information.<sup>88</sup> Because most of the general patient population see their physician on average 3.9 times a year, this creates multiple opportunities for ongoing interaction and weight management.<sup>89</sup> Moreover, Medicare, and increasingly private health insurers, cover long-term HCP counseling for obesity as long as patients lose at least 3 kg during the initial 6 months of counseling.<sup>68</sup>

### CONCLUSION

The complex, multifaceted nature of obesity presents a myriad of challenges to the treating physician. Evidence suggests that, in many cases, people with obesity are not engaged by their physician and the recommended clinical pathway is not being followed. The reasons for lack of engagement are numerous, not least the feeling that obesity treatment is futile. However, where effective strategies are implemented, the health benefits are clearly evident, even with relatively small weight loss. Using simple tools and strategies, and in some cases a multidisciplinary approach,

patients can be engaged and ongoing support and motivation may be provided, thereby improving patient care.

## ACKNOWLEDGMENTS

I am grateful to AXON Communications for medical writing and editorial assistance in the development of the submitted manuscript.

**Abbreviations and Acronyms:** **5As** = Ask, Assess, Advise, Agree, and Assist; **BMI** = body mass index; **EHR** = electronic health record; **HCP** = health care professional; **NAMCS** = National Ambulatory Medical Care Survey; **NHANES** = National Health and Nutrition Examination Survey; **NHLBI** = National Heart, Lung, and Blood Institute; **PCP** = primary care physician; **USPSTF** = US Preventive Services Task Force

**Potential Competing Interests:** Medical writing assistance was funded by Novo Nordisk A/S. Novo Nordisk was also provided with the opportunity to perform a medical accuracy review.

**Correspondence:** Address to Scott I. Kahan, MD, MPH, 1020 19th St NW, Ste 450, Washington, DC 20036 (kahan@jhu.edu).

## REFERENCES

- NCD Risk Factor Collaboration (NCD-RisC). Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet*. 2016; 387(10026):1377-1396.
- Flegal KM, Kruszon-Moran D, Carroll MD, Fryar CD, Ogden CL. Trends in obesity among adults in the United States, 2005 to 2014. *JAMA*. 2016;315(21):2284-2291.
- Sturm R, Hattori A. Morbid obesity rates continue to rise rapidly in the United States. *Int J Obes (Lond)*. 2013;37(6): 889-891.
- US Food and Drug Administration. Guidance for industry developing products for weight management. US Food and Drug Administration website. <https://www.fda.gov/downloads/Drugs/Guidances/ucm071612.pdf>. Published February 2007. Accessed June 20, 2017.
- American Medical Association House of Delegates. Resolution 420 (A-13): Recognition of obesity as a disease. May 16, 2013.
- NIH Obesity Research Task Force. *Strategic Plan for NIH Obesity Research*. Bethesda, MD: US Dept of Health and Human Services. NIH Publication No. 11-5493.
- Frellick M. AMA declares obesity a disease. Medscape website. <http://www.medscape.com/viewarticle/806566>. Published June 19, 2013. Accessed June 20, 2017.
- World Health Organization. *Obesity: Preventing and Managing the Global Epidemic: Report of a WHO Consultation*. Geneva, Switzerland: World Health Organization; 2000. WHO Technical Report Series 894.
- Yuen MS, Lui DT, Kaplan LM, et al. A systematic review and evaluation of current evidence reveals 195 obesity-associated disorders (OBAD). *Obesity Week*. 2016. Poster T-P-3166.
- Must A, Spadano J, Coakley EH, Field AE, Colditz G, Dietz WH. The disease burden associated with overweight and obesity. *JAMA*. 1999;282(16):1523-1529.
- Grover SA, Kaouache M, Rempel P, et al. Years of life lost and healthy life-years lost from diabetes and cardiovascular disease in overweight and obese people: a modelling study. *Lancet Diabetes Endocrinol*. 2015;3(2):114-122.
- Kolotkin RL, Andersen JR. A systematic review of reviews: exploring the relationship between obesity, weight loss, and health-related quality of life. *Clin Obes*. 2017;7(5):273-289.
- Withrow D, Alter DA. The economic burden of obesity worldwide: a systematic review of the direct costs of obesity. *Obes Rev*. 2011;12(2):131-141.
- Ferguson C, David S, Divine L, et al. Obesity drug outcome measures: a consensus report of considerations regarding pharmacologic intervention: George Washington University School of Public Health and Services.
- Blackman A, Foster GD, Zammit G, et al. Effect of liraglutide 3.0 mg in individuals with obesity and moderate or severe obstructive sleep apnea: the SCALE sleep apnea randomized clinical trial. *Int J Obes (Lond)*. 2016;40:1310-1319.
- Davies MJ, Bergenstal R, Bode B, et al. Efficacy of liraglutide for weight loss among patients with type 2 diabetes: the SCALE diabetes randomized clinical trial. *JAMA*. 2015;314:687-699.
- Kolotkin RL, Fujioka K, Wolden ML, Brett JH, Bjorner JB. Improvements in health-related quality of life with liraglutide 3.0 mg compared with placebo in weight management. *Clin Obes*. 2016;6:233-242.
- Li G, Zhang P, Wang J, et al. Cardiovascular mortality, all-cause mortality, and diabetes incidence after lifestyle intervention for people with impaired glucose tolerance in the Da Qing Diabetes Prevention Study: a 23-year follow-up study. *Lancet Diabetes Endocrinol*. 2014;2:474-480.
- Masmiquel L, Leiter LA, Vidal J, et al. LEADER 5: prevalence and cardiometabolic impact of obesity in cardiovascular high-risk patients with type 2 diabetes mellitus: baseline global data from the LEADER trial. *Cardiovasc Diabetol*. 2016;15:29.
- National Heart, Lung, and Blood Institute. Managing overweight and obesity in adults: systematic evidence review from the Obesity Expert Panel 2013. National Heart, Lung, and Blood Institute website. <https://www.nhlbi.nih.gov/sites/www.nhlbi.nih.gov/files/obesity-evidence-review.pdf>. Accessed June 20, 2016.
- Pi-Sunyer X, Astrup A, Fujioka K, et al. A randomized, controlled trial of 3.0 mg of liraglutide in weight management. *N Engl J Med*. 2015;373:11-22.
- Kassirer JP, Angell M. Losing weight—an ill-fated New Year's resolution [editorial]. *N Engl J Med*. 1998;338(1):52-54.
- Mann T, Tomiyama AJ, Westling E, Lew AM, Samuels B, Chatman J. Medicare's search for effective obesity treatments: diets are not the answer. *Am Psychol*. 2007;62(3):220-233.
- Look AHEAD Research Group. Eight-year weight losses with an intensive lifestyle intervention: the look AHEAD study. *Obesity (Silver Spring)*. 2014;22(1):5-13.
- Leblanc ES, O'Connor E, Whitlock EP, Patnode CD, Kapka T. Effectiveness of primary care-relevant treatments for obesity in adults: a systematic evidence review for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2011;155(7):434-447.
- Nutrition Obesity Research Centers. New insights into Americans' perceptions and misperceptions of obesity treatments, and the struggles many face. <http://www.norc.org/Research/Projects/Pages/the-asmbnrc-obesity-poll.aspx2016>; Available at <http://www.norc.org/Research/Projects/Pages/the-asmbnrc-obesity-poll.aspx>. Accessed June 19, 2016.
- Galuska DA, Will JC, Serdula MK, Ford ES. Are healthcare professionals advising obese patients to lose weight? *JAMA*. 1999; 282(16):1576-1578.
- Post RE, Mainous AG III, Gregorie SH, Knoll ME, Diaz VA, Saxena SK. The influence of physician acknowledgment of patients' weight status on patient perceptions of overweight and obesity in the United States. *Arch Intern Med*. 2011;171(4):316-321.
- Bardia A, Holtan SG, Slezak JM, Thompson WG. Diagnosis of obesity by primary care physicians and impact on obesity management. *Mayo Clin Proc*. 2007;82(8):927-932.

30. Pool AC, Kraschnewski JL, Cover LA, et al. The impact of physician weight discussion on weight loss in US adults. *Obes Res Clin Pract.* 2014;8(2):e131-e139.
31. Moyer VA. U.S. Preventive Services Task Force. Screening for and management of obesity in adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med.* 2012;157(5):373-378.
32. Jensen MD, Ryan DH, Apovian CM, et al; American College of Cardiology/American Heart Association Task Force on Practice Guidelines; Obesity Society. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society [published correction appears in *J Am Coll Cardiol.* 2014; 63(25 Pt B):3029-3030]. *J Am Coll Cardiol.* 2014;63(25 pt B): 2985-3023.
33. Physician Quality Reporting System. Measure #128: Preventive care and screening: body mass index (BMI) screening and follow-up. Vol 2017. <http://healthmonix.com/mips-pro/mips-quality-measures-2018/>. Accessed January 31, 2018.
34. Centers for Medicare & Medicaid Services (CMS), HHS. Medicare program; Merit-Based Incentive Payment System (MIPS) and Alternative Payment Model (APM) Incentive Under the Physician Fee Schedule, and Criteria for Physician-Focused Payment Models. Final rule with comment period. *Fed Regist.* 2016; 81(214):77008-77831.
35. Waring ME, Roberts MB, Parker DR, Eaton CB. Documentation and management of overweight and obesity in primary care. *J Am Board Fam Med.* 2009;22(5):544-552.
36. Crawford AG, Cote C, Couto J, et al. Prevalence of obesity, type II diabetes mellitus, hyperlipidemia, and hypertension in the United States: findings from the GE Centricity Electronic Medical Record database. *Popul Health Manag.* 2010;13(3):151-161.
37. Jackson SE, Wardle J, Johnson F, Finer N, Beeken RJ. The impact of a health professional recommendation on weight loss attempts in overweight and obese British adults: a cross-sectional analysis. *BMJ Open.* 2013;3(11):e003693.
38. Rose SA, Poynter PS, Anderson JW, Noar SM, Conigliaro J. Physician weight loss advice and patient weight loss behavior change: a literature review and meta-analysis of survey data. *Int J Obes (Lond).* 2013;37(1):118-128.
39. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for intensive behavioral therapy for obesity (210.12). Centers for Medicare & Medicaid Services website. [https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCAId=258&NcaName=Bariatric+Surgery+for+the+Treatment+of+Morbid+Obesity&ExpandComments=n&CommentPeriod=0&CALd=150&CalName=Blood+Counts+\(Addition+of+ICD-9-CM+Codes+V77.1%2C+V81.0%2C+V81.1%2C+and+V81.2+to+the+list+of+Codes+that+Do+Not+Support+Medical+Necessity\)&NCDId=353&ncover=1&CoverageSelection=National&list\\_type=ncd&bc=gAAAABAIAAAAA%3D%3D&](https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCAId=258&NcaName=Bariatric+Surgery+for+the+Treatment+of+Morbid+Obesity&ExpandComments=n&CommentPeriod=0&CALd=150&CalName=Blood+Counts+(Addition+of+ICD-9-CM+Codes+V77.1%2C+V81.0%2C+V81.1%2C+and+V81.2+to+the+list+of+Codes+that+Do+Not+Support+Medical+Necessity)&NCDId=353&ncover=1&CoverageSelection=National&list_type=ncd&bc=gAAAABAIAAAAA%3D%3D&) Updated November 2011. Accessed June 20, 2017.
40. Smith AW, Borowski LA, Liu B, et al. U.S. primary care physicians' diet-, physical activity-, and weight-related care of adult patients. *Am J Prev Med.* 2011;41(1):33-42.
41. Kraschnewski JL, Sciamanna CN, Pollak KI, Stuckey HL, Sherwood NE. The epidemiology of weight counseling for adults in the United States: a case of positive deviance. *Int J Obes (Lond).* 2013;37(5):751-753.
42. Talwalkar A, McCarty F. Characteristics of physician office visits for obesity by adults aged 20 and over: United States, 2012. *NCHS Data Brief.* 2016;(237):1-8.
43. McAlpine DD, Wilson AR. Trends in obesity-related counseling in primary care: 1995-2004. *Med Care.* 2007;45(4):322-329.
44. Yates EA, Macpherson AK, Kuk JL. Secular trends in the diagnosis and treatment of obesity among US adults in the primary care setting. *Obesity (Silver Spring).* 2012;20(9): 1909-1914.
45. Antognoli EL, Smith KJ, Mason MJ, et al. Direct observation of weight counselling in primary care: alignment with clinical guidelines. *Clin Obes.* 2014;4(2):69-76.
46. Greiner KA, Born W, Hall S, Hou Q, Kimminau KS, Ahluwalia JS. Discussing weight with obese primary care patients: physician and patient perceptions. *J Gen Intern Med.* 2008;23(5):581-587.
47. Wilkinson ML, Brown AL, Poston WS, Haddock CK, Jahnke SA, Day RS. Physician weight recommendations for overweight and obese firefighters, United States, 2011-2012. *Prev Chronic Dis.* 2014;11:E116.
48. Lewis KH, Gudzone KA, Fischer H, Yamamoto A, Young DR. Racial and ethnic minority patients report different weight-related care experiences than non-Hispanic Whites. *Prev Med Rep.* 2016;4:296-302.
49. Adams KM, Kohlmeier M, Zeisel SH. Nutrition education in U.S. medical schools: latest update of a national survey. *Acad Med.* 2010;85(9):1537-1542.
50. Kushner RF, Butsch WS, Kahan S, Machineni S, Cook S, Aronne LJ. Obesity coverage on medical licensing examinations in the United States: what is being tested? *Teach Learn Med.* 2017;29(2):123-128.
51. Gunther S, Guo F, Sinfield P, Rogers S, Baker R. Barriers and enablers to managing obesity in general practice: a practical approach for use in implementation activities. *Qual Prim Care.* 2012;20(2):93-103.
52. Huang J, Yu H, Marin E, Brock S, Carden D, Davis T. Physicians' weight loss counseling in two public hospital primary care clinics. *Acad Med.* 2004;79(2):156-161.
53. Petrin C, Kahan S, Turner M, Gallagher C, Dietz WH. Current attitudes and practices of obesity counselling by health care providers. *Obes Res Clin Pract.* 2017;11(3):352-359.
54. Ayres CG, Griffith HM. Perceived barriers to and facilitators of the implementation of priority clinical preventive services guidelines. *Am J Manag Care.* 2007;13(3):150-155.
55. Kaplan L, Golden A, Jinnett K, et al. Divergence in perceptions and attitudes among people with obesity, healthcare professionals and employers creates barriers to effective obesity management: results of the national ACTION study. Oral abstract presented at: 34th Annual Meeting of the Obesity Society; October 31-November 4, 2016; New Orleans, Louisiana. Oral abstract T-OR-2088.
56. Kolasa KM, Rickett K. Barriers to providing nutrition counseling cited by physicians: a survey of primary care practitioners. *Nutr Clin Pract.* 2010;25(5):502-509.
57. Ahmed SM, Lemkau JP, Birt SL. Toward sensitive treatment of obese patients. *Fam Pract Manag.* 2002;9(1):25-28.
58. Dietz WH, Kahan S, Gallagher C, et al. Why weight? a guide to discussing obesity and health with your patients. Strategies to Overcome and Prevent (STOP) Obesity Alliance website. <http://stopobesityalliance.org/wp-content/themes/stopobesityalliance/pdfs/STOP-Provider-Discussion-Tool.pdf>. Published 2014. Accessed June 20, 2017.
59. Vine M, Hargreaves MB, Briefel RR, Orfield C. Expanding the role of primary care in the prevention and treatment of childhood obesity: a review of clinic- and community-based recommendations and interventions. *J Obes.* 2013;2013: 172035.
60. Huizinga MM, Cooper LA, Bleich SN, Clark JM, Beach MC. Physician respect for patients with obesity. *J Gen Intern Med.* 2009;24(11):1236-1239.
61. Richard P, Ferguson C, Lara AS, Leonard J, Younis M. Disparities in physician-patient communication by obesity status. *Inquiry.* 2014;51.
62. Bravender T, Tulsky JA, Farrell D, et al. Teen CHAT: development and utilization of a web-based intervention to improve physician communication with adolescents about healthy weight. *Patient Educ Couns.* 2013;93(3):525-531.
63. Puhl R, Peterson JL, Luedicke J. Motivating or stigmatizing? public perceptions of weight-related language used by health



- providers [published correction appears in *Int J Obes (Lond)*. 2013;37(4):623]. *Int J Obes (Lond)*. 2013;37(4):612-619.
64. Puhl RM, Phelan SM, Nadglowski J, Kyle TK. Overcoming weight bias in the management of patients with diabetes and obesity. *Clin Diabetes*. 2016;34(1):44-50.
  65. Kyle TK, Puhl RM. Putting people first in obesity. *Obesity (Silver Spring)*. 2014;22(5):1211.
  66. Wittert GA, Huang KC, Heilbronn LK. Supporting the callout for people first language in obesity. *Obes Res Clin Pract*. 2015; 9(4):309.
  67. Forhan M, Risdon C, Solomon P. Contributors to patient engagement in primary health care: perceptions of patients with obesity. *Prim Health Care Res Dev*. 2013;14(4):367-372.
  68. Tang JW, Kushner RF, Cameron KA, Hicks B, Cooper AJ, Baker DW. Electronic tools to assist with identification and counseling for overweight patients: a randomized controlled trial. *J Gen Intern Med*. 2012;27(8):933-939.
  69. Canadian Obesity Network. 5As of obesity management. Canadian Obesity Network website. [http://www.obesitynetwork.ca/5As\\_adult](http://www.obesitynetwork.ca/5As_adult). Accessed November 28, 2017.
  70. Vallis M. Behaviour change counselling—how do I know if I am doing it well? the development of the Behaviour Change Counselling Scale (BCCS). *Can J Diabetes*. 2013;37(1):18-26.
  71. Rueda-Clausen CF, Benterud E, Bond T, Olszowka R, Vallis MT, Sharma AM. Effect of implementing the 5As of obesity management framework on provider-patient interactions in primary care. *Clin Obes*. 2014;4(1):39-44.
  72. Jay MR, Gillespie CC, Schlair SL, et al. The impact of primary care resident physician training on patient weight loss at 12 months. *Obesity (Silver Spring)*. 2013;21(1):45-50.
  73. Jay M, Gillespie C, Schlair S, Sherman S, Kalet A. Physicians' use of the 5As in counseling obese patients: is the quality of counseling associated with patients' motivation and intention to lose weight? *BMC Health Serv Res*. 2010;10:159.
  74. Gudzone KA, Bennett WL, Cooper LA, Bleich SN. Perceived judgment about weight can negatively influence weight loss: a cross-sectional study of overweight and obese patients. *Prev Med*. 2014;62:103-107.
  75. Aveyard P, Lewis A, Teame S, et al. Screening and brief intervention for obesity in primary care: a parallel, two-arm, randomised trial. *Lancet*. 2016;388(10059):2492-2500.
  76. Bays HE, Seger JC, Primack C, et al. Obesity Algorithm, presented by the Obesity Medicine Association. Obesity Medicine Association website. <https://obesitymedicine.org/obesity-algorithm/>. Accessed June 20, 2017.
  77. SCOPE. SCOPE certified health professional list 2017. <https://obesitymedicine.org/obesity-algorithm/>. Accessed June 20, 2017.
  78. Kushner RF, Brittan D, Cleek J, et al; ABOM Board of Directors. The American Board of Obesity Medicine: five-year report. *Obesity (Silver Spring)*. 2017;25(6):982-984.
  79. Reading JM, Snell M, LaRose JG. A systematic review of weight-related communication training for physicians: what do we know and how can we inform future development of training programs? Poster presented at: 4th Biennial D. C. Health Communication Conference; April 28-29, 2017; Fairfax, VA.
  80. Montesi L, El Ghoch M, Brodosi L, Calugi S, Marchesini G, Dalle Grave R. Long-term weight loss maintenance for obesity: a multidisciplinary approach. *Diabetes Metab Syndr Obes*. 2016; 9:37-46.
  81. Wadden TA, Butryn ML, Hong PS, Tsai AG. Behavioral treatment of obesity in patients encountered in primary care settings: a systematic review. *JAMA*. 2014;312(17):1779-1791.
  82. Welty FK, Nasca MM, Lew NS, Gregoire S, Ruan Y. Effect of onsite dietitian counseling on weight loss and lipid levels in an outpatient physician office. *Am J Cardiol*. 2007;100(1): 73-75.
  83. Dietz WH, Solomon LS, Pronk N, et al. An integrated framework for the prevention and treatment of obesity and its related chronic diseases. *Health Aff (Millwood)*. 2015;34(9): 1456-1463.
  84. Centers for Medicare & Medicaid Services. Decision memo for intensive behavioral therapy for obesity (CAG-00423N). Centers for Medicare & Medicaid Services website. <https://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?&NcaName=Intensive%20Behavioral%20Therapy%20for%20Obesity&bc=ACAAAAAIAAA&NCAId=253>. Accessed June 20, 2017.
  85. Batsis JA, Bynum JP. Uptake of the centers for Medicare and Medicaid Obesity Benefit: 2012-2013. *Obesity (Silver Spring)*. 2016;24(9):1983-1988.
  86. MacLean PS, Wing RR, Davidson T, et al. NIH working group report: innovative research to improve maintenance of weight loss. *Obesity (Silver Spring)*. 2015;23(1):7-15.
  87. Ahem AL, Wheeler GM, Aveyard P, et al. Extended and standard duration weight-loss programme referrals for adults in primary care (WRAP): a randomised controlled trial. *Lancet*. 2017; 389(10085):2214-2225.
  88. Hardcastle S, Taylor A, Bailey M, Castle R. A randomised controlled trial on the effectiveness of a primary health care based counselling intervention on physical activity, diet and CHD risk factors. *Patient Educ Couns*. 2008;70(1):31-39.
  89. O'Hara B, Caswell K. Health status, health insurance, and medical services utilization: 2010. *Curr Popul Rep*. 2013:1-16.