

9. Klompas M, Calandra T, Singer M. Antibiotics for sepsis—finding the equilibrium. *JAMA*. 2018;320(14):1433-1434.
10. Bone RC, Balk RA, Cerra FB, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Chest*. 1992;101(6):1644-1655.
11. Members of the American College of Chest Physicians & Society of Critical Care Medicine Consensus Conference Committee. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Crit Care Med*. 1992;20(6):864-874.
12. Rivers E, Nguyen B, Havstad S, et al. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med*. 2001;345(19):1368-1377.
13. Levy MM, Evans LE, Rhodes A. The Surviving Sepsis Campaign bundle: 2018 update. *Intensive Care Med*. 2018;44(6):925-928.
14. Singer M, Deutschman CS, Seymour CW, et al. The Third International Consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA*. 2016;315(8):801-810.
15. Vincent JL, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. *Intensive Care Med*. 1996;22(7):707-710.
16. Hwang SY, Jo IJ, Lee SU, et al. Low accuracy of positive qSOFA criteria for predicting 28-day mortality in critically ill septic patients during the early period after emergency department presentation. *Ann Emerg Med*. 2018;71(1):1-9.e2.
17. Usman OA, Usman AA, Ward MA. Comparison of SIRS, qSOFA, and NEWS for the early identification of sepsis in the emergency department. *Am J Emerg Med*. 2019;37(8):1490-1497.
18. Sterling SA, Puskarich MA, Glass AF, et al. The impact of the Sepsis-3 septic shock definition on previously defined septic shock patients. *Crit Care Med*. 2017;45(9):1436-1442.
19. Fang X, Wang Z, Yang J, et al. Clinical evaluation of Sepsis-1 and Sepsis-3 in the ICU. *Chest*. 2018;153(5):1169-1176.
20. Braun D. A retrospective review of the sepsis definition after publication of Sepsis-3. *Am J Med*. 2019;132(3):382-384.
21. Levy MM, Rhodes A, Phillips GS, et al. Surviving Sepsis Campaign: association between performance metrics and outcomes in a 7.5-year study. *Intensive Care Med*. 2014;40(11):1623-1633.
22. The ProCESS Investigators. A randomized trial of protocol-based care for early septic shock. *N Engl J Med*. 2014;370(18):1683-1693.
23. The ARISE Investigators and the ANZICS Clinical Trials Group. Goal-directed resuscitation for patients with early septic shock. *N Engl J Med*. 2014;371(16):1496-1506.
24. Mouncey PR, Osborn TM, Power GS, et al. ProMISe trial investigators. Trial of early, goal-directed resuscitation for septic shock. *N Engl J Med*. 2015;372(14):1301-1311.
25. Angus DC, Barnato AE, Bell D, et al. A systematic review and meta-analysis of early goal-directed therapy for septic shock: the ARISE, ProCESS and ProMISe investigators. *Intensive Care Med*. 2015;41(9):1549-1560.
26. Rhodes A, Evans LE, Alhazzani W, et al. Surviving Sepsis Campaign: international guidelines for management of sepsis and septic shock: 2016. *Crit Care Med*. 2017;45(3):486-552.
27. Monnet X, Teboul J-L. Passive leg raising: five rules, not a drop of fluid! *Crit Care*. 2015;19(1):18.
28. Miller RR, Dong L, Nelson NC, et al. Multicenter implementation of a severe sepsis and septic shock treatment bundle. *Am J Respir Crit Care Med*. 2013;188(1):77-82.
29. Mukherjee V, Evans L. Implementation of the Surviving Sepsis Campaign guidelines. *Curr Opin Crit Care*. 2017;23(5):412-416.
30. Motzkus CA, Lilly CM. Accountability for sepsis treatment: the SEP-1 core measure. *Chest*. 2017;151(5):955-957. doi:10.1016/j.chest.2017.01.011.
31. Albritton N, Omerhodzic D, Dickerson B. SEP-1 early management bundle, severe sepsis/septic shock: v5.5a measure updates and v5.0b through v5.2b analysis results. Centers for Medicare & Medicaid Services webinar 2018. www.qualityreportingcenter.com/en/inpatient-quality-reporting-programs/hospital-inpatient-quality-reporting-iqr-program/archived-events/iqr121118. Accessed April 13, 2020.
32. Aaronson EL, Filbin MR, Brown DE, et al. New mandated Centers for Medicare and Medicaid Services requirements for sepsis reporting: caution from the field. *J Emerg Med*. 2017;52(1):109-116.
33. Levy MM, Pronovost PJ, Dellinger RP, et al. Sepsis change bundles: converting guidelines into meaningful change in behavior and clinical outcome. *Crit Care Med*. 2004;32(11 suppl):S595-S597.
34. LaRosa JA, Ahmad N, Feinberg M, et al. The use of an early alert system to improve compliance with sepsis bundles and to assess impact on mortality. *Crit Care Res Pract*. 2012;2012:980369.
35. Harrison AM, Gajic O, Pickering BW, Herasevich V. Development and implementation of sepsis alert systems. *Clin Chest Med*. 2016; 37(2):219-229.
36. Iwashyna TJ, Odden A, Rohde J, et al. Identifying patients with severe sepsis using administrative claims: patient-level validation of the Angus implementation of the international consensus conference definition of severe sepsis. *Med Care*. 2014;52(6):e39-e43.



**Interested in writing
for JAAPA?**

Visit www.jaapa.com, choose the Authors and Info tab, and click on Information for Authors and Reviewers.