



## CME POST-TEST

All post-tests must be completed and submitted online.

EXPIRATION DATE: JULY 2021

Earn Category I CME Credit by reading both CME articles in this issue, reviewing the post-test, then taking the online test at <http://cme.aapa.org>. Successful completion is defined as a cumulative score of at least 70% correct. This material has been reviewed and is approved for 1 hour of clinical Category I (Preapproved) CME credit by the AAPA.

### SEPSIS

1. Which of the following is considered one of the SIRS diagnostic criteria?
  - a. temperature less than 38° C
  - b. heart rate greater than 90 beats/minute
  - c. altered mental status
  - d. end-organ damage
2. Which of the following is a component of the qSOFA score?
  - a. WBC count greater than 12,000 cells/mm<sup>3</sup>
  - b. temperature greater than 38° C
  - c. GCS score less than 15
  - d. Respirations less than 20
3. Which diagnostic test is part of the SSC 1-hour bundle?
  - a. high sensitivity troponin
  - b. CBC count
  - c. blood cultures
  - d. erythrocyte sedimentation rate
4. Which treatment is part of the SSC 1-hour bundle?
  - a. broad-spectrum antibiotics
  - b. stress-dose corticosteroids
  - c. packed red blood cell transfusion
  - d. therapeutic hypothermia
5. According to the SSC 1-hour bundle, vasopressors should be administered to achieve what BP goal?
  - a. greater than 90 mm Hg systolic
  - b. greater than 65 mm Hg MAP
  - c. greater than 85 mm Hg MAP
  - d. greater than 120 mm Hg systolic

### KAWASAKI DISEASE

6. Kawasaki disease is most commonly diagnosed in patients of which age group?
  - a. under age 5 years
  - b. age 5 to 10 years
  - c. age 10 to 16 years
  - d. age 18 to 24 years
7. Which of the following appear to decrease the chances of acquiring Kawasaki disease?
  - a. being of Northeast Asian descent
  - b. being male
  - c. breastfeeding
  - d. taking vitamin D supplements
8. Which of the following is a common clinical sign of Kawasaki disease?
  - a. currant-jelly stools
  - b. meningismus
  - c. conjunctivitis sparing the iris
  - d. supraclavicular lymphadenopathy
9. Which of the following is considered a mainstay of treatment for Kawasaki disease?
  - a. low-dose aspirin
  - b. IV immunoglobulin
  - c. broad-spectrum antibiotics
  - d. high-dose corticosteroids
10. Which of the following is a common complication of Kawasaki disease?
  - a. coronary artery aneurysm
  - b. sepsis
  - c. peripheral neuropathy
  - d. renal failure