




Hemocult Classics: The Hospitalist's Management of Acute GI Bleeding

Amanda Stark, PA-C

Mayo Clinic Arizona

October 1, 2021





Disclosures


- No relevant commercial relationships to disclose.



Learning Objectives

- At the conclusion of this session, participants should be able to:
 - Develop differential diagnosis for a patient presenting with hematemesis, hematochezia or melena
 - Discuss initial stabilization and resuscitative measures for a patient presenting with a GI bleed
 - Recognize indications for different endoscopic and radiologic diagnostic studies for localizing and treating source of GI bleeding
 - Discuss pharmacologic management of acute GI bleeds as well as secondary prophylaxis

Case #1: Mr. A



67 y/o male presents with 3-day history of melena and several weeks of intermittent dull epigastric pain and nausea.

- PMH/PSH: HTN, HLD, osteoarthritis, s/p cholecystectomy
- Medications: lisinopril, simvastatin, meloxicam, baby aspirin
- Vitals: T 36.8C, HR 102, BP 97/62, RR 22, SpO2 99%
- Exam: Pale, no jaundice, tachycardic, mild epigastric tenderness without guarding, melena in rectal vault, guaiac (+)
- Labs: Hgb 8.3, HCT 24.8, WBC 8.0, PLT 299, INR 1.0, BUN 39, Cr 1.1, remainder of CMP WNL, lipase 29



Which of the following is recommended as part of this patient's pre-endoscopic management?

- A. Transfuse 1 unit PRBCs
- B. Octreotide 50 mcg bolus + 50 mcg/hr drip
- C. Erythromycin 250 mg IV
- D. Nasogastric lavage



Upper Gastrointestinal Bleeding (UGIB)

- Bleeding from the esophagus, stomach or duodenum
 - Proximal to the Ligament of Treitz
- Typical presentation with hematemesis (bloody or coffee ground) and/or melena
 - Hematochezia in major/brisk bleeds, usually associated with hemodynamic instability
- Accounts for over half a million hospital admissions in the US annually



Initial Evaluation and Resuscitation in UGIB

- ABCs, Close monitoring – Be aware of potential for rapid decompensation
 - Tracheal intubation for airway protection if severe hematemesis or AMS
- Focused history & physical
- Labs including CBC, serial H&H, BUN/Cr, coagulation studies, type & screen
- Adequate venous access
- Fluid resuscitation, transfusion
 - Lack of clear guidelines regarding colloid vs crystalloids and rate of resuscitation
- Gastroenterology consult
- Nasogastric lavage not routinely recommended



Transfusion

- Guidelines recommend restrictive PRBC transfusion policy
 - Transfuse for hemoglobin < 7 g/dL
 - Higher for hypotensive patients due to expected equilibration after volume repletion with crystalloids
 - Threshold of hemoglobin < 8 g/dL is reasonable in patients with pre-existing cardiovascular disease



Pre-Endoscopic Medical Therapy

- Proton pump inhibitors
 - Pantoprazole or Esomeprazole 80 mg IV followed by 8mg/hr infusion, or 40 mg IV Q12 hours
 - Should be considered to reduce higher risk stigmata of hemorrhage requiring endoscopic therapy
 - Recommended if endoscopy will be delayed
 - Discontinue after endoscopy unless ulcer or erosion identified
- Prokinetics
 - Erythromycin 250 mg IV over 20-30 min; 30-90 minutes pre-endoscopy
 - May improve visualization at endoscopy → increased diagnostic yield
- Vasoactive medications if variceal bleed suspected




Management of Coagulopathy and Antithrombotic Medications in UGIB

- Anticoagulation status should not delay endoscopic intervention!
- Coagulopathy due to comorbid condition
 - Platelet transfusion thresholds not clear in guidelines – consider for PLT $<50 \times 10^9/L$
 - Correction of INR with FFP for coagulopathy of liver disease not recommended
- Antiplatelet agents
 - Reinstatement once hemostasis achieved – may require multidisciplinary decision regarding timing of resumption
 - Consider uninterrupted continuation of aspirin if indication is secondary prevention
- Anticoagulants
 - Consider reversal – Prothrombin Complex Concentrates (PCC), Fresh frozen plasma (FFP), Vitamin K
 - Risk/Benefit analysis for timing of resumption – reason for anticoagulation, etiology of bleed
- Tranexamic acid not recommended


Upper Endoscopy

- Gold standard for diagnosis and treatment of upper GI bleed.
- Recommended within 24 hours of presentation, after hemodynamic resuscitation
- Endoscopic treatment modalities – choice depends on lesion characteristics, operator experience, availability
 - Hemostatic clip
 - Sclerosant injection (absolute ethanol)
 - Thermal coagulation
 - Argon Plasma coagulation (APC) – non-contact
 - Epinephrine as combination therapy
 - Hemospray powder



Non-Endoscopic Diagnostics and Interventions for UGIB


- CT Angiography to detect active bleeding
- Transcatheter arterial embolization
 - Generally performed by IR after failure of endoscopic therapy and medical management
 - Access via common femoral artery, identification of active bleeding by contrast extravasation and selective embolization
 - Clips may be placed during EGD to help with localization
- Surgery
 - Rarely required, usually reserved for perforation or malignancy





Differential for UGIB: Peptic Ulcer Disease

Duodenal ulcer with visible vessel



Duodenal ulcer in a patient with recent upper gastrointestinal bleeding. The ulcer base (arrows) is visible as the whitish rim underlying the protruding vessel. The erythematous mound in the center of the ulcer represents an arteriole that has eroded into the lumen of the duodenum.

Courtesy of Eric D Libby, MD.

UpToDate®


- Peptic Ulcer Disease is the most common cause of UGIB
- Findings on endoscopy graded based on Forrest classification: guides endoscopic therapy & intensity of post-endoscopic PPI
 - High-risk stigmata after endoscopic hemostasis: IV PPI (either intermittent or continuous infusion) x72 hours
 - Low-risk stigmata: do not require endoscopic treatment and patient can generally be discharged after endoscopy on oral PPI therapy
- H. Pylori testing should be performed & treatment initiated if (+)
 - Very low rebleeding rates after eradication
- NSAIDs should be discontinued

Stigmata of hemorrhage	Forrest Classification
Active spurting bleeding	IA
Active oozing bleeding	IB
Non-bleeding visible vessel	IIA
Adherent clot	IIB
Flat pigmented spot	IIC
Clean base	III

Differential for UGIB: Inflammatory/Erosive Processes

- Esophagitis
 - Risk factors: history of GERD, medications (NSAIDs, PO bisphosphonates, tetracyclines), infections (*Candida*, HSV)
 - Hematemesis more common than with other bleeding causes
 - Treatment with acid suppression and risk factor reduction; endoscopic treatment rarely required
- Gastritis/Duodenitis
 - Risk factors: similar to PUD, alcohol consumption, bariatric surgery, certain autoimmune diseases
 - Often noted on endoscopy, but rarely cause significant UGIB unless coagulopathic
 - Treatment with acid suppression, occasionally endoscopic therapy with APC
- Cameron Lesions – erosions or ulcers in the sac of a hiatal hernia
 - Acute bleeding treated endoscopically
 - Consider surgical repair of hiatal hernia if recurrent bleeding

Severe esophagitis



Endoscopic view of severe reflux esophagitis in a patient who presented with upper gastrointestinal bleeding.

Differential for UGIB: Vascular Lesions

- Angiodysplasia – aka arteriovenous malformation (AVM), angiectasia, vascular ectasia
 - More common in patients >60 y/o
 - Increased incidence in ESRD, von Willebrand disease, aortic stenosis
 - Can be found throughout the GI tract
 - Bleeding tends to be recurrent/chronic, can usually be treated endoscopically
- Gastric antral vascular ectasia (GAVE)
 - Can be isolated or associated with portal hypertension or systemic sclerosis
 - Usually causes low-grade bleeding/IDA, rarely presents with acute bleed
 - Endoscopic coagulation may be used
- Dieulafoy's lesion: aberrant submucosal vessel eroding epithelium in absence of ulcer
 - Etiology unclear; Bleeding episodes often self-limited, but can be recurrent and profuse
 - Endoscopy best performed during acute bleeding, variety of hemostatic techniques used

Gastric antral vascular ectasia (watermelon stomach)



Endoscopy in a patient with gastric antral vascular ectasia (GAVE) shows the antrum and the pylorus (center) with erythematous radial stripes resembling the rind of a watermelon. The patient presented with iron deficiency anemia. Courtesy of Laurence Bailen, MD.




Differential for Upper GI Bleed: Neoplasms

- Benign or malignant
 - Esophageal
 - Gastric
- Irregular ulcer margins may suggest malignancy
- Bleeding from diffuse mucosal ulceration or from erosion into vessel
- Severe bleeding indicates poor prognosis
- Endoscopic treatments including standard hemostatic techniques, Hemospray powder may be effective

Differential for Upper GI Bleed: Traumatic/Iatrogenic Causes

- Mallory-Weiss syndrome – mucosal lacerations secondary to forceful retching
 - Usually self-limited but may see massive hemorrhage
 - Endoscopy with various hemostatic techniques used
- Foreign Body
- Post-surgical, Post-polypectomy, Post-ERCP
- Aortoenteric Fistula
 - Rare, usually iatrogenic, most common in duodenum
 - Presents with “herald bleed” followed by massive bleeding
 - High mortality rate
 - CTA for diagnosis, emergent surgical repair

Mallory-Weiss tear



Endoscopy shows a mucosal tear extending into the muscularis mucosa at the distal esophagus.

Courtesy of Moises Guelrud, MD.



Mr. A

- Mr. A received IV fluid resuscitation with improvement in HR & BP
- Hemoglobin nadired at 7.5, did not require transfusion
- ASA held since indication was primary prophylaxis
- Started on IV PPI, received erythromycin, EGD that afternoon showed ulcer with non-bleeding visible vessel in the gastric fundus treated with hemostatic clip
- Continued on IV PPI x 72 hours, d/c home with PO PPI BID
- Instructed to discontinue Meloxicam and avoid other NSAIDs

American College of Gastroenterology Clinical Guideline 2021




Figure 2. Initial management of patients presenting with overt upper gastrointestinal bleeding. ^aFuture risk assessment tools may be used if score discriminates risk of transfusion, hemostatic intervention or death with 99–100% sensitivity (0%–1% false negatives). RBC, red blood cell.

Case #2: Ms. B

53 y/o female presents with 2 episodes of coffee-ground emesis followed by 1 episode of bright red hematemesis after arrival to the ED

- PMH/PSH: T2DM, Alcohol use disorder, Cirrhosis, grade 1 varices on EGD 2 years ago, has not followed up since
- Medications: Metformin BID, Can't remember the rest, hasn't filled prescriptions recently
- Vitals: T 37.1 C, HR 115, BP 91/58, RR 27, SpO2 95%
- Exam: Mildly jaundiced, tachycardic, distended abdomen with fluid wave, slightly tremulous, no asterixis
- Labs: Hgb 7.1, HCT 21.8, WBC 11.4, PLT 94, INR 1.7, BUN 41, Cr 0.8, Na 131, K 3.2, CO2 18, Tbili 3.1, AST 311, ALT 210, lipase 137



Which medication is this patient missing that is indicated for primary prophylaxis of hemorrhage in her case?

- A. Pantoprazole
- B. Metoprolol
- C. Spironolactone
- D. Nadolol




Pre-Endoscopic Management of Suspected Variceal UGIB

- Resuscitative measures including IV access, IVF, transfusion, airway protection, correction of coagulopathy as in any UGIB
 - Avoid volume overload: risk of rebound portal hypertension → rebleeding
- Antibiotic prophylaxis
 - Ceftriaxone 1g IV daily x 7 days, can transition to PO quinolone if discharged before 7 days
 - Patients with cirrhosis and bleed should continue antibiotics for up to seven days regardless of the bleeding source
- Vasoactive medications – Octreotide 50 mcg bolus followed by 50 mcg/hr x 2-5 days
 - Splanchnic vasoconstriction → decrease portal pressure
- Hepatic encephalopathy management if present concurrently
- Prokinetics, EGD within 12 hours of presentation for suspected acute variceal hemorrhage

Differentials for UGIB in Patients with Cirrhosis

Normal Venous Anatomy

Esophageal Varices and Portal Hypertension



- Esophageal Varices
 - EGD gold standard for diagnosis & grading; routine EGDs in compensated cirrhosis, repeat if decompensation
 - Non-selective beta blockers & endoscopic band ligation for primary & secondary prevention of bleeding
 - Band ligation is treatment of choice with possible sclerotherapy to help control active bleeding
 - Stents can be used for tamponade of uncontrolled bleeding
 - Surveillance endoscopy 1-2 weeks after EV hemorrhage, TIPS recommended for recurrent bleeding
- Gastric Varices
 - Less prevalent & bleed less often than esophageal, but often more severe & less responsive to therapy so higher mortality
 - Endoscopic band ligation or cyanoacrylate injection, consider early TIPS, may require surgery
- Portal Hypertensive Gastropathy
 - Often diagnosed incidentally on endoscopy; bleeding typically chronic
 - Limited effective endoscopic options – APC or Hemospray
 - Nonselective beta blocker, Consider TIPS



Ms. B

- Received hemodynamic resuscitation with IVF, required 2 units PRBCs
- Started on Ceftriaxone and Octreotide drip
- Urgent endoscopy showed 1 large esophageal varix with stigmata of recent bleeding, 2 other medium-sized nonbleeding varices – band ligation performed
- US showed mild ascites, not enough fluid for paracentesis
- Continued on IV antibiotics & octreotide drip x3 days, Discharged on ciprofloxacin to complete 1 week, New beta blocker prescription
- Social work provided resources for IOP
- Follow up scheduled with Hepatology in 1 week




Fig. 7. Band ligation of a distal EV. (A) Acutely. (B) Follow-up band ligation.

Case #3: Mr. C

73 y/o male presents with 4 episodes of red/maroon hematochezia this week, similar episodes last month that resolved after a couple days, no pain

- PMH/PSH: Atrial fibrillation, HTN, T2DM, diastolic heart failure, CKD 3 GERD, s/p segmental resection for diverticulitis 20 years ago
- Medications: apixaban, furosemide, metoprolol, insulin, p.r.n. famotidine
- Vitals: T 36.5 C, HR 82, BP 110/84, RR 18, SpO2 98%
- Exam: No acute distress, heart irregularly irregular, no abdominal tenderness, bright red blood on rectal exam
- Labs: Hgb 9.8, HCT 32.1, WBC 7.1, PLT 180, INR 1.4, BUN 28, Cr 1.8, remainder of CMP WNL




What is an appropriate next step in management?

- A. Administer prothrombin complex concentrate (PCC)
- B. Nasogastric lavage to rule out brisk upper GI bleed
- C. Start antibiotics
- D. Order bowel prep to start tonight



Lower Gastrointestinal Bleeding (LGIB)

- Bleeding from the colon or rectum
- Accounts for approximately 20% of GI bleed cases
- Typical presentation with hematochezia
 - Rarely, right-sided colonic bleeding can present with melena
- Most stop spontaneously and have favorable outcomes



Initial Stabilization and Resuscitation

- Focused H&P and lab evaluation
- Pre-colonoscopy medical management
 - Hemodynamic resuscitation, transfusion and management of coagulopathy/antithrombotic medications as in upper GI bleeds
- Gastroenterology consult
- Exclusion of massive UGIB with EGD if hematochezia with hemodynamic instability
 - Consider nasogastric lavage if moderate suspicion for upper GI source



Colonoscopy and Flexible Sigmoidoscopy

- Colonoscopy should be initial diagnostic procedure for most patients with LGIB
 - Within 24 hours for patients with high-risk clinical features, evidence of ongoing bleeding
 - Examine entirety of colon, may intubate terminal ileum to rule out proximal blood suggestive of small bowel source
 - Push enteroscopy allows visualization of proximal 60 cm of jejunum
 - Bowel preparation – 4-6L PEG-based solution administered over 3-4 hours until clear rectal output
 - Endoscopic hemostasis if high-risk stigmata of bleeding
 - Options include clips, APC, thermal endotherapy +/- epinephrine injection
- Flexible Sigmoidoscopy
 - Examines distal 60 cm of colon, does not require full prep
 - Occasionally used as first investigation in patients with minimal bright red blood per rectum



Non-Endoscopic Diagnostics and Interventions for LGIB

- CT Angiography
 - Highly accurate at localizing site if active bleeding
 - Reasonable screening test before angiography or emergent surgery
- Tagged RBC scintigraphy
 - More sensitive for bleeding than CTA but less expedient & less accurate at localizing site → most suitable for intermittent bleeding
 - Can perform repeated scans after initial injection of tagged cells
- Angiography for selective embolization
 - Requires bleeding rate of 0.5-1.0 mL/min for site to be visualized
 - Typically reserved for cases when endoscopic treatment not feasible due to severe bleeding and/or hemodynamic compromise
- Surgical consultation for patients with high-risk clinical features and ongoing brisk bleeding



Differential for Lower GI Bleed: Diverticular

- Most common cause of LGIB
 - Prevalence increases with age
 - Most frequently located in descending and sigmoid colon, but ascending colon diverticula have higher tendency to bleed
 - Self-limited in 70-80% of cases, may be massive and life-threatening
 - Usually painless
- Endoscopic treatment for identified bleeding source generally effective with low risk of rebleeding

Differential for Lower GI Bleed: Vascular Lesions

- Angiodysplasia/Angiectasias
 - Common in right colon, prevalence increases with age
 - Can occur secondary to radiation proctitis
 - Usually present with occult bleeding
 - Usually episodic and self limited if overt
 - Venous in origin so not likely to be massive
 - Thermal endoscopic therapies (usually APC) effective, but rebleeding may occur

Endoscopic image of colonic angiodysplasia



Angiodysplasia appears endoscopically as peripherally expanding dilated capillaries with a central origin measuring between 0.1 to 1.0 cm in diameter.

Courtesy of Rome Jutabha, MD.



Differential for Lower GI Bleed: Inflammatory

- Inflammatory Bowel Disease
 - Hematochezia more common in UC than Crohn's
 - Flexible sigmoidoscopy may be indicated in active UC to assess severity & exclude infection but full colonoscopy avoided due to risk of perforation
- Infectious Colitis
 - Can generally be distinguished from other sources of bleeding by history, identification of pathogen on cultures/PCR panel
- Ischemic Colitis
 - More common in older adults, tends to be continuous and left-sided
 - Mucosal friability, often clear demarcation between involved and normal mucosa
 - Bleeding generally self limited



Differential for Lower GI Bleed: Malignancy

- Responsible for approximately 10% of cases of rectal bleeding in patients >50 y/o
- Bleeding typically low grade, recurrent
 - Melena vs hematochezia depending on site of bleeding
 - Often occult
- Endoscopic therapy limited due to friability of lesions
 - Hemospray sometimes effective



Differential for Lower GI Bleed: Anorectal Disorders

- Hemorrhoids
 - Painless hematochezia from rupture of internal hemorrhoids
 - Significant bleeding uncommon
- Anal Fissures
- Rectal Varices
 - Prevalence varies between 40-55% in cirrhotic patients, increased in patients with portal vein obstruction
 - Clinically severe bleeding uncommon
 - Endoscopic band ligation recommended
- Dieulafoy's lesion
 - More common in proximal stomach but can be found in all areas of GI tract including colon



Differential for Lower GI Bleed: Iatrogenic


- Post-surgical
- Post-polypectomy
 - Often self-limited – if no evidence of ongoing bleeding, generally safe to defer repeat colonoscopy and monitor
 - Can usually be managed with endoscopic therapy (clips or thermal hemostasis)
- Radiation Proctitis
 - Can be acute or chronic
 - Bleeding uncommon
 - Sucralfate enemas can be helpful
 - Endoscopic therapy with APC for severe symptoms not responsive to conservative management



Mr. C

- Apixaban and diuretic held
- Serial H&H remained stable
- Completed bowel prep, colonoscopy revealed diverticulum in the ascending colon with visible vessel, treated with thermal coagulation + epinephrine injection
- Discharged that afternoon, home medications resumed

Blood vessel within a colonic diverticulum



Endoscopy showing a blood vessel within a diverticulum. The blood vessel is separated from the bowel lumen only by mucosa. Over time, the vessel wall is exposed to injury along its luminal aspect, possibly leading to segmental weakness which predisposes to rupture into the lumen.

Courtesy of James B McGee, MD.

American College of Gastroenterology Clinical Guideline 2016




Figure 1. Algorithm for the management of patients presenting with acute LGIB stratified by bleeding severity. CTA, computed tomographic angiography; DAPT, dual antiplatelet therapy; EGD, esophagogastroduodenoscopy; LGIB, lower gastrointestinal bleeding; NGT, nasogastric tube; PEG, polyethylene glycol; UGIB, upper gastrointestinal bleeding.

Case #4: Mrs. D

68 y/o female presents with dyspnea on exertion and near-syncope, intermittent melena, had a negative EGD and colonoscopy last month

- PMH/PSH: HTN, moderate aortic stenosis, CAD without intervention, iron deficiency anemia, CKD 4
- Medications: amlodipine, ezetimibe, iron supplements (ASA on hold recently)
- Vitals: T 36.7 C, HR 91, BP 115/82, RR 18, SpO2 98%
- Exam: Pale, no acute distress, systolic murmur, no abdominal tenderness, dark stool guaiac positive stool on rectal exam
- Labs: Hgb 7.1, HCT 23.1, WBC 4.1, PLT 180, INR 1.0, BUN 31, Cr 2.0, Tbili 0.3, remainder of CMP WNL



What is the recommended next step in evaluation for suspected small bowel bleeding?

- A. Video Capsule Endoscopy
- B. Second look EGD
- C. CT Angiogram
- D. Tagged RBC Scintigraphy



Suspected Small Bowel Bleeding

- Uncommon, 5-10% of patients presenting with GI bleed
- Obscure GI bleed
 - Terminology previously used for bleed with no source identified on upper and lower endoscopic evaluation, now proposed to only be used after thorough examination of small bowel as well
 - “Suspected small bowel bleeding” used instead
- Overt (hematemesis, hematochezia or melena) versus Occult (positive fecal occult blood test +/- iron deficiency anemia)



Small Bowel Diagnostics

- Repeat endoscopy or colonoscopy
 - Second look examinations should be considered if patient presents with recurrent overt bleeding, or if initial exam incomplete
 - Can include push enteroscopy on EGD to examine distal duodenum/proximal jejunum or intubation of terminal ileum on colonoscopy
- Video capsule endoscopy (VCE)
 - First-line procedure for small bowel evaluation
 - Allows evaluation of entire small bowel in 79-90% of cases
 - Lack of therapeutic capabilities, but can direct interventions
- Deep enteroscopy (double-balloon or single-balloon enteroscopy, spiral enteroscopy)
 - Availability and endoscopist experience varies, techniques based on different designs of overtubes that fit over enteroscope
 - Allow for both evaluation and therapeutic intervention
- Intraoperative Enteroscopy – evaluation of small bowel at laparotomy
 - Reserved for recurrent severe bleeds with comprehensive negative evaluation, or patients in whom deep enteroscopy cannot be performed



Small Bowel Diagnostics

- Radiographic imaging
 - CT enterography
 - Higher sensitivity for detection of small bowel masses including mural-based masses
 - Consider before VCE if established IBD diagnosis, prior radiation, small bowel surgery or suspected stenosis
 - Requires ability to drink oral contrast
 - CT angiography
 - May be used for overt bleeding, can detect bleeding rates as slow as 0.3 mL/min
 - MR enterography
 - Can be considered if contraindication to CT or to avoid radiation exposure
 - Radionuclide scanning
 - Can detect lower rates of bleeding, can perform delayed imaging after injection, but less accurate localization
 - Meckel's scan – specifically for detection of Meckel's diverticulum, used in young patients
- Angiography - Not usually helpful unless significant active bleeding (requires rate of 0.5-1.0 mL/min)



Differential for Small Bowel Bleeding


- Prevalence of different causes is age dependent
- Vascular Lesions (angiectasias)
 - Most common cause of small bowel bleeding
 - Risk factors include advanced age, chronic renal failure, aortic stenosis, hereditary disorders
 - Endoscopic therapy
- Inflammatory bowel disease
- Meckel's diverticulum – most common congenital abnormality of GI tract
 - Meckel's scan and/or endoscopic evaluation, refer to surgery for consideration of resection
- Erosive Lesions (Dieulafoy, NSAID ulcers)
- Neoplasm – GIST, lymphoma, adenocarcinoma, carcinoid
- Small bowel varices or portal hypertensive enteropathy



Case #4

- Mrs. D underwent VCE with findings of angiodysplasia in the ileum with evidence of bleeding
- Subsequent double-balloon enteroscopy with thermal coagulation
- Received 1 unit PRBCs given Hgb <8 with symptoms and h/o CAD
- ASA resumed
- Discharged home, follow up with GI and Cardiology to evaluate for need for aortic stenosis

Bleeding in the ileum



Capsule endoscopy image showing bleeding in the ileum.



Take-Home Points

- Initial ABCs and hemodynamic resuscitation for any GI bleed includes adequate venous access and conservative transfusion threshold unless massive active bleeding
- Management of anticoagulation and antiplatelets on case-by-case basis, should resume once hemostasis achieved as soon as safe
- Medical management may include PPI for UGIB, octreotide and antibiotics in setting of cirrhosis
- Endoscopic management generally preferred, variety of therapy techniques available
- Multiple options for radiologic evaluation
- Consider angiography with selective embolization for brisk active bleeds
- Obscure GI bleeding terminology only used after comprehensive small bowel evaluation



References

- UpToDate
- Barkun, A. N., et al. (2019). Management of Nonvariceal upper GASTROINTESTINAL Bleeding: Guideline recommendations from the international consensus group. *Annals of Internal Medicine*, 171(11), 805. <https://doi.org/10.7326/m19-1795>
- Gerson, L. B., Fidler, J. L., Cave, D. R., & Leighton, J. A. (2015). ACG clinical guideline: Diagnosis and management of small bowel bleeding. *American Journal of Gastroenterology*, 110(9), 1265–1287. <https://doi.org/10.1038/ajg.2015.246>
- Gralnek, I. M., et al. (2021). Endoscopic diagnosis and management OF Nonvariceal upper gastrointestinal hemorrhage (nvugih): European Society of Gastrointestinal Endoscopy (ESGE) Guideline – Update 2021. *Endoscopy*, 53(03), 300–332. <https://doi.org/10.1055/a-1369-5274>
- Kovacs, T. O. G., & Jensen, D. M. (2019). Varices. *Clinics in Liver Disease*, 23(4), 625–642. <https://doi.org/10.1016/j.cld.2019.07.005>
- Laine, L., & Jensen, D. M. (2012). Management of patients with ulcer bleeding. *American Journal of Gastroenterology*, 107(3), 345–360. <https://doi.org/10.1038/ajg.2011.480>
- Laine, L., Barkun, A. N., Saltzman, J. R., Martel, M., & Leontiadis, G. I. (2021). ACG clinical Guideline: Upper gastrointestinal AND ulcer bleeding. *American Journal of Gastroenterology*, 116(5), 899–917. <https://doi.org/10.14309/ajg.0000000000001245>
- Lewis, M. (2008). Bleeding colonic Diverticula. *Journal of Clinical Gastroenterology*, 42(10), 1156–1158. <https://doi.org/10.1097/mcg.0b013e3181862ad1>
- Nett, A., & Binmoeller, K. F. (2019). Endoscopic management of Portal Hypertension–related Bleeding. *Gastrointestinal Endoscopy Clinics of North America*, 29(2), 321–337. <https://doi.org/10.1016/j.giec.2018.12.006>
- Samuel, R., Bilal, M., Tayyem, O., & Guturu, P. (2018). Evaluation and management OF non-variceal upper gastrointestinal bleeding. *Disease-a-Month*, 64(7), 333–343. <https://doi.org/10.1016/j.disamonth.2018.02.003>
- Stanley, A. J., & Laine, L. (2019). Management of acute upper gastrointestinal bleeding. *BMJ*, l536. <https://doi.org/10.1136/bmj.l536>
- Strate, L. L., & Gralnek, I. M. (2016). ACG clinical guideline: Management of patients with acute lower gastrointestinal bleeding. *American Journal of Gastroenterology*, 111(4), 459–474. <https://doi.org/10.1038/ajg.2016.41>



Questions?

Stark.Amanda@mayo.edu