

Reducing MRSA and Healthcare Associated Infections

Paul Szczybor PA-C DFAAPA
Lifebridge Critical Care

The Scourge of MDROs

- Methicillin Resistant Staph aureus (MRSA)
- Healthcare Associated Infections (HAI)
- Central Line Blood Stream Infections (CLABSI)
- Surgical Site Infections (SSI)
- Ventilator Associated Pneumonia (VAP)

Healthcare Associated Infections

- A Leading Cause of Preventable Illness and Death
- 1 in 31 Hospitalized Patients
- 70,000 Deaths per year
- ↑ LOS, ↑ Cost, ↑ Morbidity/ Mortality

Targeted Screening & Isolation



Contact Precautions

- Effective?
- Longer LOS
- ↓ Healthcare worker interaction
- ↑ Falls and pressure ulcers
- Negative Psychological effects
- Worse Patient Satisfaction Scores
- Higher 30-day Readmission Rates

Decolonization

- Chlorhexidine gluconate (CHG) bathing




Nasal Decolonization

- Mupirocin
- Povidone Iodine
- Alcohol-based sanitizer

Strategy

MRSA Screening
Contact Precautions
Targeted decolonization



Universal Decolonization

CHG bathing qday

+

Nasal sanitizer BID
(alcohol-based)

Results

- Eliminate Contact Precautions
- Decrease Isolation days
- Decrease SSI 50-81%
- Decrease MRSA bacteremia (up to 100%)
- Saved \$\$

Keys to Success

- No isolation MRSA colonized
- Universal decolonization
- Hand Hygiene
- CHG bathing daily + alcohol-based nasal sanitizer BID

Citations

- Kaur DC, Narayan PA. Mupirocin resistance in nasal carriage of *Staphylococcus aureus* among healthcare workers of a tertiary care rural hospital. *Indian J Crit Care Med* 2014;18:716-21.
- Huang SS, Septimus E, Kleinman K, Moody J, Hickok J, Avery TR, et al. Targeted versus universal decolonization to prevent ICU infections. *N Engl J Med* 2013; 368: 2255-65.
- Tran K., Bell C., Stall N., Tomlinson G., McGeer A., Morris A., Gardam M., Abrams H.B. The Effect of Hospital Isolation Precautions on Patient Outcomes and Cost of Care: A Multi-Site, Retrospective, Propensity Score-Matched Cohort Study. *J. Gen. Intern. Med.* 2017;32:262–268.

Citations

- Morgan DJ, Zhan M, Goto M, Franciscus C, Alexander B, Vaughan-Sarrazin M, et al. The effectiveness of contact precautions on methicillin-resistant *Staphylococcus aureus* (MRSA) in long-term care across the United States. *Clin Infect Dis.* 2020;71(7):1676–1683.
- Mullen A, Wieland HJ, Weiser ES, et al. Perioperative participation of orthopedic patients and surgical staff in a nasal decolonization intervention to reduce *Staphylococcus* spp surgical site infections. *AJIC.* 2017;45, 554-6. DOI: <https://doi.org/10.1016/j.ajic.2016.12.021>.
- Reeves L, Barton L, Nash M, et al, Effectiveness of an alcohol-based nasal antiseptic in reducing MRSA bacteremia in an adult intensive care population. *ICHE* 2020;41, s206. DOI: <https://doi.org/10.1017/ice.2020.748>.

Citations

- Landis-Bogush K, Belani A, Impacts of coordinated hospital-wide use of alcohol-based nasal decolonization on infection rates, patient care and cost savings. DOI: <https://doi.org/10.1016/j.ajic.2019.04.091>

Contact Information

Paul Szczybor PA-C DFAAPA

papauls@verizon.net