

LAB MEDICINE EMERGENCIES

Andy Herber, PA-C

Assistant Professor of Medicine, Mayo College of Medicine

Associate in Hospital Medicine, Mayo Clinic

CENSUS

Mr. Farmer

Ms. Colacey

Mr. Sweaty

Mrs. Sad

Mr. Etoh

MR. FARMER

PMH/PSH

Never been to doctor.

SOCIAL HISTORY:

Married. Neversmoker. No ETOH.

MEDS:

None.

***Transferred from OSF for femur fracture after falling off tractor and being rolled over.

	ADMIT LABS	DAY 2
Hemoglobin	11.7	11.1
WBC	11.2	12.7
Platelets	199,000	159,000
Creatinine	1.0	2.4
Glucose	146	189
Potassium	4.8	7.0
Sodium	144	141

WHY EMERGENCY?

- Cardiac Conduction abnormalities
- Paralysis
- Muscle Weakness

RHABDOMYOLYSIS

Traumatic	Nontraumatic Exertional	Nontraumatic Nonexertional
Crush Syndrome	Marked exertion	Drugs Toxins
Immobilization	Hyperthermia	Infections

HYPERKALEMIA

- Decreased excretion (renal dx)**
- Tissue Catabolism (rhabdo, hemolysis, GI bleed)**
- Cell shift (acidosis, lack of insulin)**
- Excessive intake (IV, PO, KCl salt substitute)**
- Blood transfusion**
- Medications (ACE/ARB, K+ sparing, cyclosporine, NSAIDS)**
- Pseudo (hemolysis, elevated WBC (> 50K) platelets (>1million))**
- Heparin Induced Hypoaldosteronism (2-4d after admin)**

HYPERKALEMIA: TREATMENT

Intervention	Mechanism	Onset and Duration	Dose	Remember
Calcium Gluconate	Stabilize cardiac membrane.	Immediate Onset, transient	1000mg (10mL of 10% solution)	Can repeat X1 if EKG changes persist. Can exacerbate dig tox.
Albuterol	Shift potassium into cells	20-30 minutes, Transient	10-20mg in 4ml nebulized solution every four hours.	Beta agonist
Insulin D50	Shift potassium into cells.	10-20minutes, transient	10units of regular insulin with D50	Monitor blood sugar closely.
Furosemide	Increases urinary potassium excretion	Onset: 5 - 30 mins Duration: 2 - 6 h	20 – 40 mg IV q12 – 24 h	Use only after hydration, Useful in volume overload, heart failure
Kayexalate	Increase potassium excretion.	1-2 hours	15-30grams orally	Do not give to post op or renal transplant pts
Dialysis	Potassium removal	Immediate	n/a	Marked tissue breakdown, ESRD on HD,
SZC/Patiromer	Cation exchange	1-4hours	10g TID x 48hrs or 8.4g daily.	?Restriction

STABILIZE

Intervention	Mechanism	Onset and Duration	Dose	Remember
Calcium Gluconate	Stabilize cardiac membrane.	Immediate Onset, transient	1000mg (10mL of 10% solution)	Can repeat X1 if EKG changes persist. Can exacerbate dig tox.

SHIFT

Intervention	Mechanism	Onset and Duration	Dose	Remember
Albuterol	Shift potassium into cells	20-30 minutes, Transient	10-20mg in 4ml nebulized solution ever four hours.	Beta agonist
Insulin D50	Shift potassium into cells.	10-20minutes, transient	10units of regular insulin with D50	Monitor blood sugar closely.

???Bicarbonate???

REMOVE

Intervention	Mechanism	Onset and Duration	Dose	Remember
Furosemide	Increases urinary potassium excretion	Onset: 5 - 30 mins Duration: 2 - 6 h	20 – 40 mg IV q12 – 24 h	Use only after hydration, Useful in volume overload, heart failure
SZC/Patiromer	Cation Exchange	1-4hours	10g TID x 48hrs	Restricted?
Dialysis	Potassium removal	Immediate	n/a	Marked tissue breakdown, ESRD on HD

MS. COLACEY

PMH/PSH

Obesity

PSH:

Right Total Knee

SOCIAL HISTORY:

Single. Wheelchair bound. Resides in SNF.

MEDS:

Fentanyl patch. MiraLAX.

ROS:

“Confusion, back pain, Can’t poop!!!”

ADMIT LABS

Lab	Admission
Hemoglobin	10.2
MCV	86
Platelets	214,000
Sodium	133
Potassium	3.9
Bicarbonate	23
Creatinine	1.5
BUN	48
Alk Phos	586
ALT	38
Bilirubin	1.0
ABG	Normal
UA	Negative

WHAT SHOULD WE ORDER NEXT?

- A. Serum Calcium**
- B. Lidoderm Patch and PT Consult**
- C. Fractionate Alkaline Phosphatase**
- D. Right Upper Quadrant US**
- E. Roseangela's Pizza**

ADMIT LABS

Lab	Admission
Hemoglobin	10.2
MCV	86
Platelets	214,000
Sodium	133
Potassium	4.8
Bicarbonate	23
Creatinine	1.5
BUN	48
Alk Phos	586
ALT	38
Bilirubin	1.0
ABG	Normal
UA	Negative
Calcium	14.6

ADMIT LABS

Lab	Admission
Hemoglobin	10.2
MCV	86
Platelets	214,000
Sodium	133

Calcium 14.6

ALT	38
Bilirubin	1.0
ABG	Normal
UA	Negative
Calcium	14.6

WHY IS THIS AN EMERGENCY?

- Coma
- Cardiac Arrhythmia
- Renal failure
- Polyuria
- Polydipsia
- Dehydration
- Nausea
- Anorexia
- Muscle Weakness
- Constipation
- Nephrolithiasis

Prostate

Breast

Kidney

Thyroid

Lung

*****multiple myeloma*****

Type	Frequency	Bone metastasis	Causal agent	Tumors
Humoral hypercalcemia of malignancy	80%	Minimal, absent	PTHRP	Squamous cell ca (head/neck, esophagus), cervix, renal, endometrial, ovarian, breast, HTLV- lymphoma
Local osteoclastic hypercalcemia	20%	Extensive, common	Cytokines, chemokines, PTHrP	Breast, multiple myeloma, lymphoma
1-25 Di-OH D secreting lymphoma	< 1 %	Variable	Vitamin D analogues	Lymphomas

What else causes hypercalcemia?

HYPERCALCEMIA

- Primary Hyperparathyroidism**
- Malignancy (PTH peptide, Bone Mets)**
- Sarcoidosis**
- Drugs (HCTZ, Lithium, Theophylline, Vitamin D)**
- Vitamin D intoxicification**
- Hyperthyroidism**
- Immobilization**

TOTAL VS IONIZED?

Corrected Calcium = Calcium + 0.8 (4.0-Albumin)

Calcium

Hyperalbuminemia

Hypoalbuminemia

Chronic Kidney Disease

Ionized

\$\$\$\$\$\$\$\$\$\$\$\$\$\$\$\$\$\$

Acid Base disorders

Hyperparathyroidism

Hyperphosphatemia

HYPERCALCEMIA: TREATMENT

Intervention	Mechanism	Duration of action	Dose	Remember
Normal saline	Restores volume, promotes calcium excretion	Hours	200-300 ml/hr IV to maintain UOP 100-150 cc/hr	Caution in heart failure
Bisphosphonates	Block osteoclast mediated bone resorption	Onset: 1 – 3 d Duration: 2 - 4 wk.	Pamidronate IV 60-90mg (2 - 4 h) Zoledronic acid IV 4 - 8 mg (15 min)	Caution in renal failure Rare: Osteonecrosis of the jaw, esp. with repeated doses
Calcitonin	Increases calcium excretion Decreases Ca reabsorption	Onset: 1 – 4 h Duration: 6 – 12 h	4 - 8 IU/Kg q12h SQ/IV	Safe Tachyphylaxis limits use
Furosemide	Increases urinary Ca excretion	Onset: 5 - 30 mins Duration: 2 - 6 h	20 – 40 mg IV q12 – 24 h	No longer 1 st line Rx, Use only after hydration, Useful in volume overload, heart failure
Corticosteroids	Inhibit cytokine mediated inflammation, vit D analog	Onset: 1-5 d Duration: 2-4 wks.	Prednisone 40-60 mg/d	Useful in lymphoma, granulomatous disease

HYDRATE!!!!

Intervention	Mechanism	Duration of action	Dose	Remember
Normal saline	Restores volume, promotes calcium excretion	Hours	200-300 ml/hr IV to maintain UOP 100-150 cc/hr	Caution in heart failure

MOVE CALCIUM

Intervention	Mechanism	Duration of action	Dose	Remember
Bisphosphonates	Block osteoclast mediated bone resorption	Onset: 1 – 3 d Duration: 2 - 4 wk.	Pamidronate IV 60-90mg (2 - 4 h) Zoledronic acid IV 4 - 8 mg (15 min)	Caution in renal failure Rare: Osteonecrosis of the jaw, esp. with repeated doses
Calcitonin	Increases calcium excretion Decreases Ca reabsorption	Onset: 1 – 4 h Duration: 6 – 12 h	4 - 8 IU/Kg q12h SQ/IV	Safe Tachyphylaxis limits use
Furosemide	Increases urinary Ca excretion	Onset: 5 - 30 mins Duration: 2 - 6 h	20 – 40 mg IV q12 – 24 h	No longer 1 st line Rx, Use only after hydration, Useful in volume overload, heart failure

Intervention	Mechanism	Duration of action	Dose	Remember
Calcitonin	Increases calcium excretion Decreases Ca reabsorption	Onset: 1 – 4 h Duration: 6 – 12 h	4 - 8 IU/Kg q12h IV	Safe Tachyphylaxis limits use

Tachyphylaxis?????????

?

TACHYPHYLAXIS:

Definition:

**rapid development of tolerance or
immunity to the effects of a drug**

OTHER

Intervention	Mechanism	Duration of action	Dose	Remember
Corticosteroids	Inhibit cytokine mediated inflammation, vit D analog	Onset: 1-5 d Duration: 2-4 wks.	Prednisone 40-60 mg/d	Useful in lymphoma, granulomatous disease
Denosumab	Rank Ligand Inhibitor	Within 24hrs	60mg subcu may repeat x1	Not renally cleared.

MS. SAD

PMH

Diabetes

PSH:

4 Bunion Removals

SOCIAL HISTORY:

Widowed. Neversmoker. Recently placed in SNF by you.

MEDS:

Lisinopril 5mg daily, ASA 325mg daily, Zoloft 100mg daily (NEW)

****Vitals stable. Discharged three weeks ago, direct admit bounceback from SNF for confusion****

LABS

Lab	Discharge	Admission
Hemoglobin	10.9	10.2
WBC	7000	8600
Platelets	186000	214,000
Sodium	136	121
Potassium	4.6	3.9
Glucose	133	146
Bicarbonate	19	23
Creatinine	0.9	1.1
BUN	19	16
AST	54	52
ALT	91	100
ABG	n/a	Normal
UA	n/a	Negative

LABS

Lab	Admission
Hemoglobin	10.2
WBC	8600
Na	121
Glucose	146
Bicarbonate	23
Creatinine	1.1
BUN	28
AST	52
ALT	100
ABG	Normal
UA	Negative

WHY IS THIS BAD?

- Cerebral Edema
- Seizures
- Coma
- Respiratory Arrest
- Malaise
- Nausea
- Headache
- Altered Mental Status

Hypovolemic

- Volume contraction
- Sweating, Diarrhea, or Vomiting
- Diuretics (Thiazides, Loop)
- Cerebral Salt Wasting

Euvolemic


- SIADH (Head trauma, Seizure, CNS disease, Neoplastic, Meds)
- Adrenal Failure
- Hypothyroidism


Hypervolemic


- Congestive Heart Failure
- Cirrhosis
- Polydipsia
- Nephrotic Syndrome
- Renal disease


Pseudo

- Hyperglycemia *FOR EVERY 100 ABOVE 100 ADD 1.6*
- Hypertriglyceridemia
- Paraproteinemia










LABS



Lab	Admission
Sodium	121
Creatinine	1.1
BUN	18
Vitals	Normotensive
Edema	None
Serum Osmo	260
Orthostatics	Normal
Urine Sodium	58
Urine Osmo	300

SIADH

- **Malignancy** (Lung, Brain, GI, GU, lymphoma)
- **Pulmonary** (Pneumonia, asthma, COPD, Lung CA)
- **Intracranial** (Trauma, Stroke, Hemorrhage, Infection)
- **Drugs** (Antipsycotics, Antidepressants, Chemo, Ectasy)
- **Misc** (Pain, Nausea, Post Operative)

DIAGNOSING SIADH

Serum Osmo <275	Normal Acid Base Status
Euvolemic	Normal Adrenal Function
Urine Osmo >100 Osm/kg	Normal renal function
Urine Sodium > 40mEq/L	Normal Thyroid

TREATMENT PEARLS

Volume Contraction = Normal Saline

Fluid Overload = Diurese and Restrict

SIADH = Fluid Restrict (800mL per day)

3% Saline = 100mL bolus (2-3meq)

Goal <9meq in 24 hours

Osmotic Demyelination Syndrome (ODS)

MR. SWEATY

PMH

Diabetes

Chronic Kidney Disease

PSH:

Appendectomy

SOCIAL HISTORY:

Single. Significant alcohol and tobacco use.

MEDS:

NPH 15u BID, ASA 325mg Qday, Metoprolol 50mg BID, Norvasc 5mg BID

ROS:

Nausea, vomiting, weight loss. Night sweats.

ADMIT LABS

Lab	Admission
Hemoglobin	8.2
WBC	1,900
Platelets	54,000
Sodium	133
Potassium	6.3
Bicarbonate	23
Creatinine	1.5
LDH	645
Phosphorus	8.6
Calcium	6.8
Uric Acid	9.6

CAIRO-BISHOP CLASSIFICATION

LABORATORY TUMOR LYSIS SYNDROME

Uric acid $\geq 8\text{mg/dL}$ ($\geq 476\mu\text{mol/L}$) or 25% increase from baseline

Phosphorus $\geq 6.5\text{mg/dL}$ ($\geq 2.1\text{mmol/L}$) or 25% increase from baseline

CLINICAL TUMOR LYSIS SYNDROME

Creatinine > 1.5 times the upper limit of normal

Cardiac arrhythmia or sudden death

Seizure

WHY EMERGENCY?

Hyperuricemia

- Uric acid can crystallize in renal tubules and lead to acute renal failure

Hyperkalemia

- Life-threatening arrhythmias

Hyperphosphatemia

- Leads to hypocalcemia, tetany, seizures, arrhythmias

ADMIT LABS

Lab	Admission
Hemoglobin	8.2
WBC	1,900
Platelets	54,000
Sodium	133
Potassium	6.3
Bicarbonate	23
Creatinine	1.5
LDH	645
Phosphorus	8.6
Calcium	6.8
Uric Acid	9.6

ADMIT LABS

Lab	Admission
Hemoglobin	8.2
WBC	1,900
Platelets	54,000
Sodium	133
Potassium	5.8
Bicarbonate	23
Creatinine	1.5
LDH	645
Phosphorus	8.6
Calcium	6.2
Uric Acid	9.6

Problem	Intervention	Dosage	Comments
Renal insufficiency	IV fluids	Normal saline, 3L/m2 daily	Caution if history of CHF
Hyperuricemia	<p>Allopurinol (xanthine oxidase inhibitor) inhibits uric acid production</p> <p>Rasburicase (recombinant urate oxidase) Converts uric acid into water soluble allantoin</p>	<p>100 mg/m2 divided in 3 doses (every 8 hours) Commonly 600 mg initially followed by 300 mg/d (prophylaxis)</p> <p>IV 0.15-0.2 mg/kg/d (can be given up to 5 days)</p>	<p>Reduce dose in renal failure. Only effective in prophylaxis. Not recommended if uric acid level above 7.5 mg/dl</p> <p>For severe hyperuricemia. Contraindicated in pregnancy, G6PD deficiency Expensive</p>
Hyperphosphatemia	<p>Phosphate binders</p> <p>Dialysis</p>	50-150 mg/kg daily	<p>Low phosphorus diet Dialysis if refractory to therapy</p>
Hyperkalemia	<p>Insulin (Regular) Dextrose (50%)</p> <p>Calcium gluconate (10%)</p> <p>Kayexalate</p> <p>Sodium bicarbonate</p> <p>Albuterol</p> <p>Dialysis</p>	<p>10 units IV 5 ml Dextrose 50% IV push</p> <p>1000 mg IV</p> <p>20 mg nebulized</p>	<p>If hyperkalemic ECG changes</p> <p>Dialysis if refractory to therapy</p>
Hypocalcemia	Calcium gluconate (10%)	1000 mg IV (no faster than 200mg IV/minute)	Use only if symptomatic, Caution with severe hyperphosphatemia

MR. ETOH

PMH

None.

PSH:

None.

SOCIAL HISTORY:

Single. Significant for LOTS of alcohol and tobacco use.

MEDS:

None.

ROS:

Unknown.

LABS

Lab	Admission
Hemoglobin	10.2
MCV	101
Platelets	14,000
Sodium	128
Potassium	2.9
Bicarbonate	33
Creatinine	1.4
BUN	28
AST	252
ALT	112
Lactate	4.6
INR	4.1

LABS

Lab	Admission
Hemoglobin	10.2
MCV	101
Platelets	14,000
Sodium	128
Potassium	2.9
Bicarbonate	33
Creatinine	1.4
BUN	28
AST	252
ALT	112
Lactate	4.6
INR	4.1

ETOH

Decreases platelet aggregation 20 minutes after ingestion

Toxic to megakaryocytes

Causes liver disease and splenomegaly

Accelerates platelet apoptosis

Decreased TPO

Extra Credit:

MCV goes Up (Folate)

AST/ALT ratio 2:1 and typically less than 300

Magnesium down

DIC

Sepsis

Malignancy (leukemia, mucinous tumors (eg, pancreatic, gastric, ovarian), and brain tumors).

Trauma (central nervous system)

Infection (Bacterial, COVID)

Obstetrical complications

Intravascular hemolysis

WHAT IS THIS DIC YOU SPEAK OF?

Disseminated Intravascular Coagulation

Coagulation and Fibrinolysis at same time

Blood exposed to procoagulant (TF, LPS)

Acute vs Chronic

Herber.Andrew@mayo.edu

REFERENCES

Clinician's Guide to Laboratory Medicine

Desai, Samir MD 2009

Henry's Clinical Diagnosis and Management by Laboratory Methods

21st Edition. McPherson, Richard, Pincus, Matthew. 2007

The Hyponatremic Patient: A systemic approach to laboratory diagnosis

Milionis, Et al.

CMAH:2002 April 16; 166 (8) 1056-1062

Oncologic Emergencies: Pathophysiology, Presentation, Diagnosis, and Treatment